



Research

Increasing uptake of hepatitis C virus infection case-finding, testing, and treatment in primary care: evaluation of the HepCATT (Hepatitis C Assessment Through to Treatment) trial

Jeremy Horwood, Clare Clement, Kirsty Roberts, Cherry-Ann Waldron, William L Irving, John Macleod and Mathew Hickman

British Journal of General Practice 2020; 70 (697): e581-e588. DOI: <https://doi.org/10.3399/bjgp20X708785>

[Article](#)[Figures & Data](#)[Info](#)[eLetters](#)[PDF](#)

Abstract

Background Hepatitis C virus (HCV) infection is a key cause of liver disease but can be cured in more than 95% of patients. Around 70 000 people in England may have undiagnosed HCV infection and many more will not have been treated. Interventions to increase case-finding in primary care are likely to be cost-effective; however, evidence of effective interventions is lacking. The Hepatitis C Assessment Through to Treatment (HepCATT) trial assessed whether a complex intervention in primary care could increase case-finding, testing, and treatment of HCV.

Aim To investigate the feasibility and acceptability of the HepCATT intervention.

Design and setting A qualitative study with primary care practice staff from practices in the south west of England taking part in the HepCATT trial.

Method Semi-structured interviews were carried out with GPs, nurses, and practice staff to ascertain their views of the HepCATT intervention at least 1 month after implementing the intervention in their practice. Normalisation process theory, which outlines the social processes involved in intervention implementation, informed thematic data analysis.

Results Participants appreciated the HepCATT intervention for increasing knowledge and awareness of HCV. Although some initial technical difficulties were reported, participants saw the benefits of using the audit tool to systematically identify patients with HCV infection risk factors and found it straightforward to use. Participants valued the opportunity to discuss HCV testing with patients, especially those who may have been previously aware of HCV risk. Future implementation should consider fully integrating software systems and additional resources to screen patient lists and conduct tests.

Conclusion When supported by a complex intervention, primary care can play a crucial role in identifying and caring for patients with HCV infection, to help stem the HCV epidemic, and prevent HCV-related illness.

[hepatitis C](#)[primary care](#)[qualitative research](#)[randomised controlled trial](#)[screening](#)

PDF

Help

INTRODUCTION

Hepatitis C (HCV) is a bloodborne virus. Acute infection is usually asymptomatic, with symptoms not appearing until after several years of chronic infection, when the liver may be severely damaged, leading to patients presenting late with cirrhosis and liver disease.¹ In the UK, injecting drug use is the main route of HCV transmission; other routes include unscreened blood transfusion, unsafe health care, exposure to blood-contaminated needles, for example, by tattooing, and sexual practices that lead to exposure to blood.^{1,2} Research from 2019 suggests that up to 43% of future hepatitis C infections could be prevented if transmission among people who inject drugs was reduced.³ Treatment of chronic HCV infection results in sustained viral clearance in a high percentage (>95%) of treated patients.⁴ An estimated 143 000 people in the UK have chronic HCV infection;^{2,5} however, HCV has been described as a silent epidemic, with <50% of those infected aware of their HCV status and with many more not receiving treatment, who are therefore at increased risk of morbidity, mortality, and onward transmission.^{2,5,6} HCV screening interventions have been effective in increasing case finding in target populations, for example, in migrants and attenders at drug treatment centres;^{7,8} however, those not in contact with specialist services are not systematically being assessed. The crucial role of primary care in stemming the HCV epidemic and preventing HCV-related illness has been highlighted,⁹ although many primary care patients at risk of HCV infection have not been tested.¹⁰

The World Health Organization (WHO) has called for an increase in HCV diagnosis, setting a target of 90% of infected people in Europe knowing their status by 2030.¹¹ To increase diagnosis and treatment, the National Institute for Health and Care Excellence (NICE) recommends implementing cost-effective interventions to increase case finding in primary care;¹² however, robust evidence is lacking, and case finding and treatment rates in many sites are low.¹³

The Hepatitis C Assessment Through to Treatment (HepCATT) trial was conducted to assess whether a complex intervention in primary care could increase the identification and treatment of HCV-infected patients compared with usual care.¹⁴ The HepCATT intervention, following NICE recommendations, consisted of training practice staff about HCV; providing patients with information about HCV; and use of an electronic patient record algorithm to identify patients with HCV infection risk factors or patients who had been previously diagnosed and not referred for treatment. During the trial, 2071 (16%) of patients identified in the intervention practices and 1163 (10%) in the control practices were tested for HCV, giving an overall intervention effect as an adjusted risk ratio of 1.59 (95% CI = 1.21 to 2.08, $P < 0.001$).¹⁵ Intervention practices had a greater yield of positive antibody tests compared with control practices (6.2% versus 4.2%; rate ratio 1.42; 95% CI = 0.95 to 2.13), and the intervention was demonstrated to be highly cost-effective for the NHS.¹⁵ Recommendations have been made by the trial team to implement the HepCATT intervention across the UK.

Although the Royal College of General Practitioners and the National Institute for Health and Care Excellence provide guidance for GPs on testing for hepatitis C, research suggests that the current practice of case-finding, testing, and treatment is not effective. This study examined the implementation of the HepCATT (Hepatitis C Assessment Through to Treatment) trial complex intervention, which consists of training practice staff; providing patient HCV posters and leaflets; and using an electronic patient record algorithm for proactive case-finding of patients with HCV infection risk factors or patients diagnosed >1 year previously with no evidence of referral for treatment. The intervention was found to be feasible and acceptable to staff, and could be effective in supporting primary care to follow hepatitis C infection testing and treatment guidelines.

How this fits in


This article describes the HepCATT trial nested qualitative study, which aimed to investigate practice staff members' views and experiences of intervention acceptability, and the facilitators and challenges to wider implementation of the intervention.

METHOD

Study design

The HepCATT trial was a pragmatic two-armed practice-level cluster randomised controlled trial in 45 primary care practices (22 intervention and 23 control) in the south west of England. Practices were randomised to either control (usual care) or intervention. The intervention comprised:

- online HCV educational resources (Royal College of General Practitioners e-learning module — Hepatitis C: Enhancing Prevention, Testing and Care) and 1-hour face-to-face staff training on the epidemiology, diagnosis, management of HCV infection, and trial processes;
- patient posters and leaflets explaining HCV risk factors and treatment options in practice waiting rooms; and
- Audit+ software (Informatica Systems Ltd) integrated into practice electronic patient record systems and used to identify patients with risk factors for HCV infection or patients who had been diagnosed >1 year previously without a referral for treatment.¹⁴



Patients identified by the audit tool were invited by letter/email (and followed up by telephone, email, or text) for an HCV test, and automatically flagged by the software by creating on-screen pop-ups to encourage opportunistic testing if the patient attended a consultation.

Semi-structured interviews were conducted with staff in intervention practices involved in implementing the intervention to investigate their views and experiences of the intervention; the acceptability and feasibility of the intervention; the impact on working practice; and attitudes to future implementation.

Sampling and recruitment

Purposive sampling was used to capture maximum variation in views and experiences, and to reflect a range of practices (in relation to size and location based on socioeconomic deprivation), and staff (in relation to professional roles). The socioeconomic status of practice populations was estimated using the Index of Multiple Deprivation decile.¹⁶ Staff from all 22 intervention practices were invited to participate in an interview after delivering the intervention for ≥ 1 month via email or telephone. Sample size was informed by the concept of 'information power',¹⁷ with analysis and sampling conducted in parallel and continuous assessment of the suitability of the information within the sample with regard to study objectives.

Data collection

Telephone Interviews were conducted by an experienced social science researcher. Audio-recorded verbal consent was gained before the interviews, which lasted between 12 and 47 minutes, with an average time of 25 minutes. A flexible topic guide was used to assist questioning but allow participants to introduce new issues (**Box 1**).

Existing HCV identification practice before HepCATT intervention

Acceptability of the intervention

Experiences of using the software

Views on opportunistic prompts

Feasibility of the intervention

Appropriateness of patients identified

Difficulties experienced with offering the test

Support needs

Approaching patients for testing opportunistically

Patient information support needs

Perceived patient views about being approached for testing

Future implementation of the intervention in primary care

Improvements to intervention

Potential barriers

Recommendations for roll-out of the intervention

PDF

Help

HCV = hepatitis C virus. HepCATT = Hepatitis C Assessment Through to Treatment

Box 1.

Interview discussion topic guide

Data analysis

Interviews were audio-recorded, transcribed, anonymised, imported to NVivo (version 10), and thematically analysed.¹⁸ Transcripts were coded inductively to establish an initial analysis framework and three researchers coded a subset of two transcripts; discrepancies were discussed to ensure a coding consensus and maximise rigour.¹⁹ The four normalisation process theory constructs²⁰ were used to further develop themes across the dataset. Normalisation process theory proposes that implementation of interventions is dependent on staff fulfilling four criteria (**Box 2**).²⁰ Emergent analysis was discussed in multidisciplinary trial management group meetings to ensure that findings were trustworthy and credible.

1. Coherence: sense making — understanding and opinion of the intervention purpose

2. Cognitive participation: buy-in — engagement with the intervention

3. Collective action: the work of putting the intervention into operation

4. Reflexive monitoring: appraisal of the intervention

Box 2.

Normalisation process theory criteria

RESULTS

Fourteen healthcare professionals (HCPs) (seven GPs, three nurses, and four practice staff) were interviewed from across 12 practices — five in areas of high deprivation (Table 1). Findings are presented for each of the normalisation process theory constructs, illustrated with anonymised verbatim quotes.

Table 1.

Participant characteristics

View inline

1. Coherence (sense making)

All HCPs described their practice of HCV testing before HepCATT as opportunistic and sporadic, testing patients with a history of drug dependency, abnormal liver function, and/or other infections such as HIV or hepatitis B. Only two HCPs described other HCV infection risk factors, such as blood transfusions given abroad, a family member with HCV, or high-risk sexual activity: *“If somebody came, you know, and we found they had HIV, we definitely tested. Um, I think if they had a hepatitis B they were tested as well but I don’t think it was routinely picked up on for anybody else.”*“(Site 17, Nurse)”

Participants understood the relevance of the intervention and valued its perceived impact. Initial engagement was led by an interest in finding out whether there were patients with HCV infection risk factors at their practice who had not yet been previously identified: *“I was interested to see whether we would have anybody, because my first thoughts were was “oh, we will be fine because you know [doctor] does it and the drugs workers do it all” ... I thought “oh, maybe we have missed people but we will wait and see”.*“(Site 6, GP)” *“I think it would be a sensible thing to do because I think that if we can diagnose people with hepatitis C earlier then potentially we can reduce their risk of becoming cirrhotic.”*“(Site 9, GP)”

2. Cognitive participation (buy-in)

Participants valued the training for enhancing HCV awareness and increasing knowledge, particularly in practices with lower levels of patients with existing HCV. Even those with previous experience of HCV believed the training had expanded their knowledge of risk factors for HCV infection and who to target for testing: *“Hepatitis C was something I was aware of anyway, perhaps more so than other GPs because of my work as a drugs lead ... certainly that was always a population that I’m aware that we should be screening ... but I think that the way HepCATT worked was potentially broadening out that ... it was a broader remit to think of hepatitis C.”*“(Site 11, GP)” *“I don’t think it was on people’s radar particularly because there’s so many other things going on ... the doctors all said “you know, we hadn’t even thought about that”*“(Site 18, Nurse)”

The training itself acted as a prompt for opportunistic testing: *“I think having the discussion at the beginning of the trial to raise awareness and remind people about the risk factors of hepatitis C probably led to some people offering the test perhaps where they wouldn’t have done .”*“(Site 11, GP)”

3. Collective action (putting HepCATT into operation)

Audit tool

Some practices encountered ‘a few teething problems’ (Site 11, GP) with setting up the audit tool, compatibility issues, and not always having the necessary time and resources; however, with support from the trial team and the software company, difficulties were resolved, and practices went on to use the audit tool without issue: *“It took a bit of getting used to but I felt like we had support to find our way around it ...*

you had to take the time just to focus on it and get used to it, but it worked fine when you were familiar with it ... that [support from trial team] helped a lot." (Site 2, Administrator)"

Participants saw the benefits of identifying patients using the audit tool and most liked that it was 'straightforward' to use and helpful in 'identifying the right people'. (Site 2, Administrator): *"I think it's fairly robust at identifying the patient that is at risk ."* (Site 9, GP)"

Screening lists of patients with HCV infection risk factors produced by the audit tool were perceived to be mostly appropriate. Although the algorithm automatically excluded certain patients (such as those on opioid patches for pain relief or who were terminally ill), issues arose in some practices with inconsistencies in medical record coding leading to inappropriate patients being identified. This was particularly pertinent to how drug users were categorised. This meant that, for some, the benefits of using the Audit+ software were offset by additional resource and time needed to screen lists. To reduce GP workload, in some practices administrators or nurses reviewed patient lists before GP screening: *"I think the only problem was about the codes that we'd used ... we had one partner in particular who tended to code this little lady who couldn't get off her sleeping tablets as drug dependent and that was a problem for us, because that's flagged up unnecessary ones ."* (Site 5, GP)" *"She [administrator] would pull off the lists and have a quick scan ... highlight the ones she thinks aren't right ... then I look through it ... so 10 sets of notes ... I would pull the notes up, quickly have a look to see if there is anything that I mean is palliative care, they have just had a consultation that hasn't been coded ."* (Site 6, GP)"

HCV testing via invite letters

Some participants were initially concerned about the resources needed to implement the intervention. Resource concerns were threefold: first, to manage the mail-out to identified patients; second, to deal with patient queries; and, third, to conduct the HCV tests. To address this, practices sent letters out in batches: *"We haven't sent out the whole lot, we would only do like 20 at a time, so we didn't want 20 people to ring up saying they wanted a test, that would have had a big impact on the practice, but it didn't turn out to be the case ."* (Site 6, GP)"

Participants also worried about disappointing patients by not being able to schedule a test promptly. They believed this would result in patients not trying to book a test at a later date. Therefore, one practice appointed a dedicated nurse to liaise with patients and conduct tests: *"We don't want them [patients] to be disappointed, because if they ring up and we say "oh, that's in a month's time" ... they won't be interested ."* (Site 6, GP)"

A few practices commented that engaging patients with a history of injecting drug use was challenging because they often did not respond to HCV testing invite letters and the practice had to 'chase up the patients' (Site 18, Nurse); however, most HCPs described that patients were happy to be approached and they did not receive any negative feedback: *"We haven't had any negative erm comments from patients saying "Why have I been picked?" or anything like that ... if the doctors have identified more risk for the letter, then obviously, I go through it with them and explain why their name has come up. And they're very understanding of why their name has been picked up ."* (Site 13, Nurse)"

A few HCPs described being cautious about approaching patients for testing with risk factors other than injecting drug use, such as organ transplants or past blood transfusions, because they did not want to scare them, but the intervention allowed for testing options to be discussed: *"But it [intervention] kind of opened up the conversation with them ... we had to be very careful, we didn't want to scare patients and the ones that probably fell into that ball park were the ones who'd had transfusions or some sort of procedure where they might be at risk but a long time ago. That was a little bit difficult, but once we had explained it to them, why it was a problem and why it might be a risk and they kind of were happy to then go ahead ."* (Site 5, GP)"

Only one practice described a negative response from a small number of HIV-positive patients, because they felt they were being stigmatised and they were already screened regularly for HCV at the local hospital: *"Some of our HIV patients, I mean two or three, were a little bit upset and also you know when I went into their records, they're all seen regularly at hospital and they do have regular Hep C tests there ."* (Site 11, Administrator)"

Opportunistic HCV testing via 'pop-up' computer screen alerts

Because of the busy nature of primary care, the computer screen alerts or pop-ups to opportunistically identify patients with HCV risk factors during routine consultations were commonly not welcomed or used by GPs. Patients often presented with complex needs and immediate clinical problems were prioritised over discussing HCV risk and testing given the limited consultation time. Participants also spoke of pop-up fatigue, of having too many screen pop-ups when dealing with a patient and therefore the impact of the alerts becoming diluted. Using the screen pop-ups was also made more challenging by the requirement for GPs to log in to the Audit+ software to activate them, which they often forgot or opted not to do: *"Just remembering to switch it on in the morning was probably the biggest issue ... but then I suspect that probably*



GPs because of the number of pop-ups it fired at you would have ignored it or try to switch it off at some point .""(Site 11, GP)" "We recognised that people were just a bit swamped and couldn't factor that into their 10-minute appointment .""(Site 5, GP)"

Although GPs did not welcome the screen pop-ups, some nurses thought they could be useful to keep HCV testing on people's 'radar': "I'm always very happy with the opportunistic testing because if I've got a pop-up box it's fantastic. I will see it and speak to the patients about it .""(Site 18, Nurse)"

4. Reflexive monitoring (appraisal of HepCATT intervention)

When appraising the intervention and making recommendations for future implementation, participants suggested streamlining information technology systems. Integrating the algorithm and screen pop-up systems within existing electronic patient record systems (for example, EMIS) would remove the need for HCPs to additionally log in to the Audit+ software: "Trying to make that alert part of EMIS might be better, because having a separate bit of software ... it did rely on each doctor logging in every morning and so perhaps having a prompt in EMIS might be more reliable ... as we tend to use prompts in EMIS more .""(Site 10, GP)"

The record search algorithm was based on a large number of Read codes and risk factors were not weighted. A few participants suggested refining the search algorithm so it could be tailored by the practice to focus on more high-risk patients and reduce the number of inappropriate patients identified. This would in turn reduce practice staff workload in screening lists of patients identified and sending invites, and would allow realistic financial planning: "Being a bit more focused on the patients who are more at risk ... it felt like it was literally picking everyone up who might have the minutest risk of hepatitis ... and it didn't always feel appropriate ... maybe it felt like the search could be narrowed down a bit than perhaps using a too broad a remit .""(Site 11, GP)" "We send out a lot of letters for a minimal response ... when there's more acute patients that could be identified ... whether that's something in the audit or the coding ... that identifies the high-risk rather than medium-risk patients. ""(Site 13, Nurse)"

Some participants suggested changing the wording of the HCV testing invite letter to alleviate staff concerns regarding sending the letters to certain patient groups: "If you have, you know, something like "please don't be offended but we are just sending letters out to anybody who's had a Hep C blood test in the last 20–25 years" ... we did have to be sort of careful if somebody might have had one for fertility blood test and they maybe now had a new partner, you know, whether picked up that letter it's, you know, they might be thinking "oh, you know, why have you got this?" .""(Site 11, Administrator)"

Implementing HepCATT in more practices could be enabled by increasing activities to raise patient and HCP awareness and acceptance of HCV and testing. A public health campaign and ongoing clinician training were suggested: "My awareness will gradually fade. ""(Site 14, GP)" "I think it's the patients accepting it ... but then that's education of the public isn't it? Everybody knows if you have bleeding down below, you have to go and see your doctor, if you cough up blood, go and see your doctor as there has been big public campaigns ... but as far as I'm aware, I haven't seen them [hepatitis C campaigns].""(Site 6, GP)"

Some participants suggested additional resources would be required for practices to implement the intervention: "If you're asking more practices to do more work then practices need to be funded to do more work, you can't just expect surgeries to take on more work. You get people phoning and speaking to the receptionist about the letter they received, you get people making appointments, you have the cost of the blood test ... they might need an appointment to discuss the results .""(Site 8, GP)"

DISCUSSION

Summary

The HepCATT intervention was valued by primary care staff for enhancing systematic identification of patients with HCV infection with the potential to benefit from treatment, compared with previous opportunistic and sporadic HCV testing. HepCATT training enhanced awareness of HCV and improved knowledge of HCV infection risk factors. Although there were initial teething problems with setting up the audit tool in some practices, most staff found it straightforward and helpful, and valued having a comprehensive list of patients to target for testing. Extra resources were required to screen lists and to conduct tests, and practices often drew on the expertise of nurses to reduce GP workload. Fees for running the algorithm and contingency staff management to conduct tests should be considered in wider implementation of the intervention. Staff valued the opportunity to discuss HCV testing with patients, especially those who may not have been previously aware of HCV risk; however, practices need to be cautious not to additionally stigmatise patients with conditions such as HIV, particularly those who may receive regular screening for HCV in secondary care. Although the algorithm excluded patients who had been tested for HCV within the previous 12

PDF

Help

months, it relied on patient record data. Therefore, if a patient had recently received an HCV test in secondary care, test results may not have been apparent on patient records at the time of the record search. As with any algorithm-based intervention, the quality and completeness of medical records are critical for its effectiveness.

When considering further use of the intervention, participants suggested refining the search algorithm to weight risk factors to reduce the number of inappropriate patients identified. Participants also recommended fully integrating information technology systems, especially computer screen pop-up software with electronic patient record systems. Views on the value of screen pop-up prompts were mixed, with some GPs highlighting 'pop-up fatigue', while some nurses valued reminders to consider HCV testing. Pop-up alerts have been found to be beneficial in influencing clinicians' behaviour;²¹ however, the danger of electronic patient record-based alerts overload, producing 'pop-up fatigue', is a common phenomenon.^{22,23} Future implementation of HepCATT should ensure that on-screen pop-ups are tailored to ensure a balance between adequate alerting and burden.

Strengths and limitations

Participants were recruited using a diverse sample of practices, GPs, practice nurses, and information technology and/or administration staff to ensure that all aspects of the intervention were captured, including software installation and use, case finding, patient consultation, and testing. Analysis demonstrated a high degree of similarity between views and experiences, suggesting acceptability across practices serving different communities. The use of normalisation process theory to inform analysis allowed for examination of issues with both the intervention design and its implementation. A study limitation is that no interviews were conducted with patients, which should be taken into consideration when interpreting the results. Although HCPs did discuss how the intervention was received by patients, which provided some insight into patients' views, there were no problems raised by HCPs except for a few patients living with HIV. These patients, because of delays with previous HCV test results from secondary care, should have been excluded from screening lists.

Comparison with existing literature

HepCATT trial findings demonstrated an effective and cost-effective intervention.¹⁵ Although the effect size was modest, the intervention is very low cost to implement. Previous research has suggested that a high proportion of patients at risk of HCV are not being tested in primary care as a result of clinicians not remembering to test patients with risk factors, and not being able to quickly and reliably determine HCV risk and status from EMIS coding.¹⁰ The current study demonstrates that HepCATT can overcome these issues, with staff valuing the intervention as a straightforward means of identifying at-risk patients and providing an opportunity to discuss HCV testing. As HCV infection is usually asymptomatic, staff welcomed the HepCATT intervention proactive case finding based on HCV risk factors. The HepCATT intervention aligns with the 'new service model for the 21st century' aspiration of the *NHS Long Term Plan* to be more proactive in providing services to enable individuals 'to take more control of how they manage their physical and mental wellbeing'.²⁴ As other HCV primary care interventions have found,²⁵ the current study indicated resource concerns for proactive case finding and increasing HCV testing; however, the findings demonstrate that nurses might be considered to alleviate increased GP workload. Previous research has also identified nursing support as a critical facilitator to increasing primary care HCV testing and as effective in supporting adherence to HCV testing and treatment guidelines.²⁶

Implications for practice

With adequate resources and technology, primary care can play an important role in identifying patients with HCV infection who have the potential to benefit from treatment. While there needs to be a multipronged approach to increase HCV testing in a range of settings, including prisons and drug treatment centres, the cost-effective HepCATT intervention provides primary care with a range of tools to improve the identification and care of patients with HCV infection, and prevent HCV-related illness. This could help the UK to reach the WHO target of infected people knowing their status by 2030 and help to stem the HCV epidemic, reducing the risk of morbidity, mortality, and onward transmission.¹¹

Acknowledgments

The authors would like to thank the general practices and patients that participated in the trial, Informatica Systems Ltd, and the NIHR Clinical Research Network West of England, which assisted with identifying and accessing GP practices to participate in the trial. The Hepatitis C Trust provided expertise and support on hepatitis C virus prevention, and essential health education materials.

Notes

Funding

The trial (ISRCTN61788850) was funded by the National Institute for Health Research (NIHR) Health Policy Research Programme (Evaluation of interventions designed to increase diagnosis and treatment of patients with hepatitis C virus infection in primary care and drug treatment settings, 015/0309). This study was designed and delivered in collaboration with the Bristol Randomised Trials Collaboration, a UK Clinical Research Collaboration Registered Clinical Trials Unit (CTU), which, as part of the Bristol Trials Centre, is in receipt of NIHR CTU support funding. Support was given by NIHR Health Protection Research Unit (HPRU) in Evaluation of Interventions and NIHR HPRU Blood Borne & Sexually Transmitted Infections, NIHR programme Grant EPIToPe, and NIHR Applied Research Collaboration West, NIHR School for Primary Care Research, and Centre for the Development and Evaluation of Complex Interventions for Public Health Improvement. The views expressed in this article are those of the authors and not necessarily those of the NHS, NIHR, or the Department of Health and Social Care.

Ethical approval

NHS ethics approval was received from the National Research Ethics Service Committee South West-Frenchay (ref: 15/SW/0094).

Provenance

Freely submitted; externally peer reviewed.

Competing interests

The authors have declared no competing interests.

Discuss this article

Contribute and read comments about this article: bjgp.org/letters

Received October 10, 2019.

Revision requested November 7, 2019.

Accepted November 25, 2019.

©The Authors

This article is Open Access: CC BY-NC 4.0 licence (<http://creativecommons.org/licenses/by-nc/4.0/>).

REFERENCES

1. Degenhardt L, Charlson F, Stanaway J, et al. Estimating the burden of disease attributable to injecting drug use as a risk factor for HIV, hepatitis C, and hepatitis B: findings from the Global Burden of Disease Study 2013. *Lancet Infect Dis* 2016;16(12):1385-1398. [CrossRef](#) [PubMed](#) [Google Scholar](#)
2. Public Health England (PHE). *Hepatitis C in the UK 2019. Working to eliminate hepatitis C as a major public health threat*. London: PHE; 2019. [Google Scholar](#)
3. Trickey A, Fraser H, Lim AG, et al. The contribution of injection drug use to hepatitis C virus transmission globally, regionally, and at country level: a modelling study. *Lancet Gastroenterol Hepatol* 2019;4(6):435-444. [Google Scholar](#)
4. Fathi H, Clark A, Hill NR, Dusheiko GE. Effectiveness of current and future regimens for treating genotype 3 hepatitis C virus infection: a large-scale systematic review. *BMC Infect Dis* 2017;17(1):22. [CrossRef](#) [PubMed](#) [Google Scholar](#)
5. Harris RJ, Harris HE, Mandal S, et al. Monitoring the hepatitis C epidemic in England and evaluating intervention scale-up using routinely collected data. *J Viral Hepatol* 2019;26(5):415-51. [CrossRef](#) [PubMed](#) [Google Scholar](#)
6. Harris RJ, Thomas B, Griffiths J, et al. Increased uptake and new therapies are needed to avert rising hepatitis C-related end stage liver disease in England: modelling the predicted impact of treatment under different scenarios. *J Hepatol* 2014;61(3):505-37. [CrossRef](#) [PubMed](#) [Google Scholar](#)
7. Flanagan S, Kunkel J, Appleby V, et al. Case finding and therapy for chronic viral hepatitis in primary care (HepFREE): a cluster-randomised controlled trial. *Lancet Gastroenterol Hepatol* 2019;4(1):32-44. [Google Scholar](#)
8. Harrison GI, Murray K, Gore R, et al. The Hepatitis C Awareness Through to Treatment (HepCATT) study: improving the cascade of care for hepatitis C virus-infected people who inject drugs in England. *Addiction* 2019;114(6):1113-1122. [CrossRef](#) [PubMed](#) [Google Scholar](#)



9. «Budd J, Robertson R Hepatitis C and general practice: the crucial role of primary care in stemming the epidemic *Br J Gen Pract* 2005;55:1325-1326
FREE Full Text Google Scholar

10. «Datta S, Horwood J, Hickman M, Sharp D Case-finding for hepatitis C in primary care: a mixed-methods service evaluation *Br J Gen Pract* 2014;DOI: <https://doi.org/10.3399/bjgp14X677112>. Google Scholar

11. «World Health Organization (WHO) *Action plan for the health sector response to viral hepatitis in the WHO European Region* Geneva WHO 2017
Google Scholar

12. «Martin NK, Miners A, Vickerman P *Assessing the cost-effectiveness of interventions aimed at promoting and offering hepatitis C testing to injecting drug users: an economic modelling report* London NICE 2012 Google Scholar

13. «Bajis S, Dore GJ, Hajarizadeh B, et al. Interventions to enhance testing, linkage to care and treatment uptake for hepatitis C virus infection among people who inject drugs: a systematic review *Int J Drug Policy* 2017;47:344-356 CrossRef PubMed Google Scholar

14. «Roberts K, Macleod J, Metcalfe C, et al. Hepatitis C — Assessment to Treatment Trial (HepCATT) in primary care: study protocol for a cluster randomised controlled trial *Trials* 2016;17:366 CrossRef PubMed Google Scholar

15. «Roberts K, Macleod J, Metcalfe C, et al. Cost effectiveness of an intervention to increase uptake of hepatitis C virus testing and treatment (HepCATT): cluster randomised controlled trial in primary care. *BMJ* 2020;doi:10.1136/bmj.m322 Abstract/FREE Full Text Google Scholar

16. «Department for Communities and Local Government *The English Index of Multiple Deprivation (IMD) 2015 — guidance 2015* https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/464430/English_Index_of_Multiple_Deprivation_2015_-_Guidance.pdf (accessed 18 Feb 2020). Google Scholar

17. «Malterud K, Siersma VD, Guassora AD Sample size in qualitative interview studies: guided by information power *Qual Health Res* 2016;26:1317-1334
CrossRef PubMed Google Scholar

18. «Braun V, Clarke V Using thematic analysis in psychology *Qual Res Psychol* 2006;3:277-101 CrossRef Google Scholar

19. «Tracy SJ Qualitative quality: eight 'big-tent' criteria for excellent qualitative research *Qual Inq* 2010;16:1083-1091 CrossRef Google Scholar

20. «May C, Finch T Implementing, embedding, and integrating practices: an outline of normalization process theory *Sociology* 2009;43:353-355 CrossRef
Google Scholar

21. «Schedlbauer A, Prasad V, Mulvaney C, et al. What evidence supports the use of computerized alerts and prompts to improve clinicians' prescribing behavior? *J Am Med Inform Assoc* 2009;16:453-458 CrossRef PubMed Google Scholar

22. «Embi PJ, Leonard AC Evaluating alert fatigue over time to EHR-based clinical trial alerts: findings from a randomized controlled study *J Am Med Inform Assoc* 2012;19:e145-e148 CrossRef PubMed Google Scholar

23. «Ancker JS, Edwards A, Nosal S, et al. Effects of workload, work complexity, and repeated alerts on alert fatigue in a clinical decision support system *BMC Med Inform Decis Mak* 2017;17:36 CrossRef PubMed Google Scholar

24. «NHS England *NHS Long Term Plan* <https://www.england.nhs.uk/long-term-plan/> (accessed 18 Feb 2020). Google Scholar

25. «Cullen W, Stanley J, Langton D, et al. Hepatitis C infection among injecting drug users in general practice: a cluster randomised controlled trial of clinical guidelines' implementation *Br J Gen Pract* 2006;56:532-538 Abstract/FREE Full Text Google Scholar

26. «Cullen W, O'Leary M, Langton D, et al. Guidelines for the management of hepatitis C in general practice: a semi-qualitative interview survey of views regarding content and implementation *Ir J Med Sci* 2005;174:323-327 PubMed Google Scholar




▲ Back to top

◀ Previous Article

Next Article ▶

In this issue



British Journal of General Practice

Vol. 70, Issue 697

August 2020

Table of Contents

Index by author

Download PDF

Article Alerts

Email Article

Citation Tools

Share

Tweet

Mendeley

Keywords

hepatitis C, primary care, qualitative research, randomised controlled trial, screening

We recommend

Hepatitis C infection among injecting drug users in general practice: a cluster randomised controlled trial of clinical guidelines' implementation

Walter Cullen, Br J Gen Pract, 2006

General practice characteristics associated with rates of testing and detection of hepatitis C: cross-sectional study in Nottingham and Derbyshire

Carol Coupland, Br J Gen Pract, 2006

Case-finding for hepatitis C in primary care: a mixed-methods service evaluation

Shivani Datta, Br J Gen Pract, 2014

Cost effectiveness of an intervention to increase uptake of hepatitis C virus testing and treatment (HepCATT): cluster randomised controlled trial in primary care

Kirsty Roberts et al., The BMJ, 2020

Cost effectiveness of an intervention to increase uptake of hepatitis C virus testing and treatment (HepCATT): cluster randomised controlled trial in primary care

Kirsty Roberts et al., The BMJ: Research, 2020

Clinical Diabetes CME Series: Anemia in CKD in Primary Care

Stephen Brunton et al, Clinical Diabetes

Powered by **TREND MD**I consent to the use of Google Analytics and related cookies across the TrendMD network (widget, website, blog). [Learn more](#)

Yes

No

PDF

Help

▼ More in this TOC Section

Primary care transformation in Scotland: qualitative evaluation of the views of national senior stakeholders and cluster quality leads

Variation in suspected cancer referral pathways in primary care: comparative analysis across the International Benchmarking Cancer Partnership

Association between hip pain and radiographic hip osteoarthritis in primary care: the CHECK cohort

Show more Research

► Related Articles

► Cited By...

Intended for Healthcare Professionals




Tweets liked by @BJGPjournal

**Andrew Papanikitas**

@gentlemedic

Editing some speculative fiction submitted to @BJGPjournal on the @OxfordTube -author for this work has added a recommended soundtrack to her narrative too! FYI @euan_lawson @nadafkhan @drricharmitage @SamMerriel @drtomround @johnspicer3



Sep 24, 2022

[Embed](#)[View on Twitter](#)

NAVIGATE

[Home](#)<https://bjgp.org/content/70/697/e581>

12/13

[Current Issue](#)[All Issues](#)[Online First](#)[Authors & reviewers](#)

RCGP

[BJGP for RCGP members](#)[BJGP Open](#)[RCGP eLearning](#)[InnovAiT Journal](#)[Jobs and careers](#)

MY ACCOUNT

[RCGP members' login](#)[Subscriber login](#)[Activate subscription](#)[Terms and conditions](#)

NEWS AND UPDATES

[About BJGP](#)[Alerts](#)[RSS feeds](#)[Facebook](#)[Twitter](#)

AUTHORS & REVIEWERS

[Submit an article](#)[Writing for BJGP: research](#)[Writing for BJGP: other sections](#)[BJGP editorial process & policies](#)[BJGP ethical guidelines](#)[Peer review for BJGP](#)

CUSTOMER SERVICES

[Advertising](#)[Contact subscription agent](#)[Copyright](#)[Librarian information](#)

CONTRIBUTE

[BJGP Life](#)[eLetters](#)[Feedback](#)

CONTACT US

[BJGP Journal Office](#)[RCGP](#)[30 Euston Square](#)[London NW1 2FB](#)[Tel: +44 \(0\)20 3188 7400](#)[Email: \[journal@rcgp.org.uk\]\(mailto:journal@rcgp.org.uk\)](#)