



**LOTHIAN HEALTH SERVICES ARCHIVE  
EDINBURGH UNIVERSITY LIBRARY  
GEORGE SQUARE  
EDINBURGH  
EH8 9LJ**

**TEL: +44 (0)131 650 3392**

**FAX: +44 (0)131 650 6863**

**URL: <http://www.lhsa.lib.ed.ac.uk>**

**EMAIL: [lhsa@ed.ac.uk](mailto:lhsa@ed.ac.uk)**

**BOOK-SCANNED 2020**

**ACCESSION No.:** LHB26/28/3/2  
**TITLE:** World Medical Journal  
**AUTHOR:** World Medical Journal

N.B. Scanned as single pages.  
Blank pages have been omitted.  
Due to variations in the density of the text some words may be illegible.  
Some text may be illegible or missing due to hole punches/creasing/smudging.  
**THIS IS THE BEST COPY AVAILABLE.**

**© Lothian Health Services Archive  
© NHS Lothian**

This PDF is supplied for the use of evidence-gathering in the UK Infected  
Blood Inquiry 2018 only.  
Scanning and subsequent digital copies have been managed in line with  
agreed procedures on data protection and information sharing.

**Please address all enquiries to the LHSA Archivist.**

CUTTER

For shock...  
exemplary  
service in  
patient  
protection

**Plasmanate®**  
plasma protein  
fraction [Human]  
5% solution, U.S.P.



First and foremost, you can count on Plasmanate to help reverse the emergency condition of circulatory collapse. Its convenient and immediate administration normalizes plasma volume quickly. But, in addition, Plasmanate protects the patient after the operation. This well-accepted human blood protein fraction has no history of coagulation defect or hepatitis transmission. *Nor will it interfere with crossmatching if future blood transfusions follow.*

Plasmanate is a 5% solution of selected human plasma proteins with stabilizers in a buffered, balanced electrolyte solution. Contains 88% serum albumin, 7% alpha globulin, 5% beta globulin. Significant electrolytes are 110 mEq. sodium, 50 mEq. chloride and a maximum of 2 mEq. potassium per liter.

Each flask is heat-treated at 60° C. for 10 hours against the possibility of transmitting the hepatitis virus.

**Administration:** Plasmanate should be administered by intravenous route only. For full details, please examine literature.

**Precautions:** Should be administered cautiously in patients with normal or increased blood volume. Package directions contain indications and all known contraindications. In new 50 ml. vial (pediatric size) and 250 and 500 ml. flasks complete with ready-to-use administration set.

*World Leader in Human Plasma Fractions*

**CUTTER Laboratories, Inc., Berkeley, California 94710, U.S.A.**

*Other Cutter Offices:* Colombia: Apartado 8263, Bogota • Mexico: Calzada México-Tacuba Núm. 1419, México 17, D.F. • Panama: Apartado 852, Panama • Canada: 1912 Tenth Avenue, S.W., Calgary, Alberta