T: 57 P.O.M.

CANVINADOLIST Normal Immunoglobulin Injection B.P. Son in Sol and a sol and

PRECAUTION

index tendos reinal in albeir in presión and anonancina poetenaria en la construcción de la construcción de la

SHORTEA-HODIOLEGICALIONS

and a percentam every such to be received with here.

32/04-1

GAMMABULIN Is a concentrate of the antibodies present in the Immune globulin (IgG) fraction of human plasma. The antibody content in GAMMABULIN Is ten to twenty times that of normal human plasma. It is

nanufactured from pooled plasma.

All plasma units are exclusively obtained from licensed plasmapheresis centers.

for the manufacture of GAMMABULIN only plasma units are used which are non-reactive in tests for HBs-antigen and HIVantibodies.

CANNABULIN is proufactured by subjecting this Souce) to a cold ethanch fractionation procedure according to Cont HIV which was added in defined quantities to samples drawn in the individual steps of the procedure was shown to be partitioned or inactivated, which led to a cumulative decrease in virus titre of at least 21 logs. The use of Innuno's cold ethanol fractionation procedure in the manufacture of, and many years of clinical experience with immunoglobulins, in which no case of anti-Him seroconversion or viral hepatitis has been reported make the transmission of viruses causing these is those through this product highly unlikely.

In the manufacture of GAHMABULIN, the immunoglobulins in their native form remain unchanged. The biological halflife and the functions depending on intact antibody molecules remain unimpaired.

GAMMABULIN is tested for identity, sterility, freedom from pyrogens, safety and antibody content.

. 1