



1 16 November 2018  
2 EMA/CHMP/BWP/303353/2010 Rev 3  
3 Committee for Medicinal Products for Human Use (CHMP)

4 **CHMP position statement on Creutzfeldt-Jakob disease**  
5 **and plasma-derived and urine-derived medicinal products**  
6

Draft Agreed by Biologics Working Party	12 September 2018
Adoption by CHMP for release for consultation	18 October 2018
End of consultation (deadline for comments)	31 October 2019

7  
8 This CHMP position statement replaces the CHMP position statement on Creutzfeldt-Jakob disease and  
9 plasma-derived and urine-derived medicinal products (EMA/CHMP/BWP/303353/2010).

Comments should be provided using this template. The completed comments form should be sent  
to [BWPsecretariat@EMA.GRO-C](mailto:BWPsecretariat@EMA.GRO-C)

10

<b>Keywords</b>	Creutzfeldt-Jakob disease, human Transmissible Spongiform Encephalopathies, plasma-derived medicinal products, urine-derived medicinal products, sporadic CJD, genetic CJD, iatrogenic CJD, variant CJD, blood infectivity, transmissibility
-----------------	--

11  
12

13 CHMP position statement on Creutzfeldt-Jakob disease  
14 and plasma-derived and urine-derived medicinal products

15 **Table of contents**

16 **Summary ..... 3**

17 **1. Introduction ..... 4**

18 **2. Human TSEs current status ..... 6**

19 2.1. Sporadic, genetic and iatrogenic forms of human TSEs.....6

20 2.2. Variant CJD .....6

21 **3. Human tissue distribution of infectivity/abnormal prion protein..... 8**

22 **4. Infectivity in blood and transmissibility via blood ..... 8**

23 4.1. Animal blood.....8

24 4.2. Human blood ..... 10

25 **5. Detection techniques..... 11**

26 **6. Leucoreduction and specific prion affinity filters ..... 12**

27 **7. Manufacturing processes for plasma-derived medicinal products..... 14**

28 **8. Infectivity in urine..... 15**

29 8.1. Animal urine ..... 15

30 8.2. Human urine..... 15

31 **9. Recommendations and proposals ..... 16**

32 9.1. Sporadic, genetic and iatrogenic CJD and plasma-derived medicinal products ..... 16

33 9.2. Variant CJD and plasma-derived medicinal products ..... 16

34 9.2.1. Exclusion Criteria..... 17

35 9.2.2. Leucoreduction and specific prion affinity filters ..... 18

36 9.2.3. Manufacturing processes for plasma-derived medicinal products..... 19

37 9.2.4. Recall of batches where information becomes available post-donation..... 20

38 9.2.5. Albumin used as an excipient or in manufacturing processes..... 21

39 9.2.6. Substitution with alternative products ..... 21

40 9.2.7. Optimal Use..... 21

41 9.3. Urine-derived medicinal products ..... 21

42 **References ..... 23**

43

44 This is the third revision of the CHMP Position Statement on "Creutzfeldt-Jakob disease and plasma-  
45 derived and urine-derived medicinal products". It was originally published in February 2003  
46 (EMA/CPMP/BWP/2879/02), replacing the CPMP Position Statement on "New variant CJD and plasma-  
47 derived medicinal products" (CPMP/201/98) from February 1998. EMA/CPMP/BWP/2879/02 was  
48 revised in June 2004 (EMA/CPMP/BWP/2879/02 rev 1.) and in June 2011  
49 (EMA/CHMP/BWP/303353/2010).

## 50 **Summary**

51 The purpose of this revision is to account for scientific developments since the last revision in 2011.  
52 The scientific information has been updated. However, there is no change in the regulatory  
53 recommendations regarding exclusion, potential testing of donors, the need to evaluate the prion  
54 reduction capacity of the manufacturing process and batch recalls.

55 Emergence of variant CJD (vCJD) was noted in UK in 1996. Although the number of cases has been in  
56 decline in the UK since 2001, isolated cases of vCJD are still being identified since 2011 in the UK as in  
57 other countries and there is still uncertainty about the future number of cases. Studies on appendix  
58 tissues from the UK indicate a potential high prevalence (about 1:2000 in the people examined) of  
59 infected persons and this is of concern considering potential human-to-human transmissions. However,  
60 there are some uncertainties about the significance of the results and their correlation to the BSE  
61 epizootic. The recent appendix tissue studies from 2013 have not produced a clear answer to the  
62 question of whether abnormal prion in the British population is limited to those exposed to the BSE  
63 epizootic. Residence in the UK has been a recognised risk factor for vCJD and there is no change to the  
64 recommendations for country-based donor exclusion. It is recommended that donors who have spent a  
65 cumulative period of 1 year or more in the UK between the beginning of 1980 and the end of 1996 are  
66 excluded from donating blood/plasma for fractionation. In addition, there is no change in the  
67 recommendations for precautionary recall of batches of plasma-derived medicinal products where a  
68 donor to a plasma pool subsequently develops vCJD.

69 Originally, a wider distribution and higher level of infectivity or abnormal prion protein in human  
70 peripheral tissues, including the lymphoreticular system was found in patients with vCJD compared  
71 with sporadic CJD. However, recent studies indicate that the prion-levels in peripheral tissue may vary  
72 in individual vCJD patients, and some cases of vCJD and sporadic CJD have been found with equal  
73 amounts of abnormal prion protein or seeding activity in peripheral tissue. Moreover, infectivity was  
74 detected in the plasma of two in four sCJD infected patients tested by bioassay in human PrP  
75 transgenic mice. These findings raise a concern that sCJD could be present in plasma from donors  
76 incubating sCJD. However, a direct link between sCJD cases and treatment with plasma-derived  
77 medicinal products or blood has not been established and there is still no firm epidemiological evidence  
78 that sporadic, genetic or iatrogenic forms of human TSEs have been transmitted from person to person  
79 through exposure to blood, plasma products or urinary-derived medicinal products. Therefore, at this  
80 stage, the recommendation not to recall batches of plasma-derived medicinal products where a donor  
81 is later confirmed as having sporadic, genetic or iatrogenic CJD is maintained, provided the  
82 manufacturer has demonstrated using appropriate methodology, that the process includes steps which  
83 significantly minimize the risk of prion contamination of the final product.

84 No recommendation for testing of donors was made in the former version of this position statement  
85 and this policy is maintained. Significant progress has been made in developing sensitive in vitro  
86 assays for prion detection in blood and some methods offer the possibility for screening of blood  
87 donors. However, these tests have not yet been completely validated according to the current

88 requirements of specificity as defined in the Common Technical Specifications for *in vitro* diagnostics.  
89 Comparison and validation of potential screening tests has been considerably confounded by the  
90 paucity of blood samples from confirmed cases of clinical prion disease and very limited samples  
91 available from asymptomatic individuals who later developed vCJD.

92 No requirement for leucoreduction of plasma was made in the former version of this position statement  
93 and this policy is maintained. Experience in TSE animal models indicates that leucodepletion reduces  
94 the risk for transmission by blood transfusion. However with respect to plasma-derived medicinal  
95 products, the same models indicate no clear evidence that leucoreduction of plasma significantly  
96 reduces the risk of prion disease transmission.

97 Taking account of the available data concerning potential contamination of blood donations with vCJD  
98 or CJD agents, assuring an adequate prion reduction capacity of the manufacturing process is  
99 considered crucial for the TSE safety of plasma-derived medicinal products. Available data indicate that  
100 the manufacturing processes for plasma-derived medicinal products would reduce TSE-infectivity if it  
101 were present in human plasma. Manufacturers have been required to estimate the potential of their  
102 specific manufacturing processes to reduce infectivity using a step-wise approach and it has been  
103 recommended that manufacturers consult the relevant competent authorities at each of the milestones  
104 in this estimation. This policy is maintained.

105 In support of this recommendation, CHMP and BWP, with the involvement of external experts,  
106 developed guidance on how to investigate manufacturing processes with regard to vCJD risk. Since  
107 publication of this Guideline in 2004, the methods for prion detection, the knowledge about infectivity  
108 in prion area in general and, prion infectivity in the blood have significantly evolved. Experimental  
109 studies highlighted the fact that prion removal capacity may vary directly according to the spiking  
110 preparation (dispersion and TSE agents strains) particularly for steps based on retention mechanisms.

111 There is no change to the recommendations for urine-derived medicinal products. Low levels of  
112 infectious TSE agents were first detected in the urine of scrapie-infected rodents and in the urine of  
113 deer with chronic wasting disease raising concerns about the possibility of infectious agents in human  
114 urine. Recent investigations on human urine have produced diverse results. While one study failed to  
115 detect infectivity by bioassay in the urine from 3 sCJD patients using sensitive assays, abnormal prion  
116 protein has recently been detected in urine from 8 out of 20 sCJD patients, 1 of 2 iatrogenic cases as  
117 well as in 1 of 13 vCJD patient urine samples using a highly sensitive immunoassay. There is still no  
118 epidemiological evidence of CJD or vCJD transmission by urine-derived medicinal products and prion  
119 reduction capacity of the manufacturing processes has been indicated. Therefore, the recommendation  
120 to apply exclusion criteria for selection of a urine donor panel from the former version of the position  
121 statement is maintained. The same exclusion criteria should be applied with respect to sJD and vCJD  
122 as used for blood/plasma donors providing starting material for the manufacture of plasma-derived  
123 medicinal products and manufacturers should follow up these criteria at defined intervals.  
124 Manufacturers of urine-derived medicinal products are recommended to estimate the potential of their  
125 manufacturing processes to reduce infectivity by following a similar general stepwise approach as  
126 recommended for plasma-derived medicinal products.

## 127 **1. Introduction**

128 Creutzfeldt-Jakob disease (CJD) is a rare neurodegenerative disease belonging to the group of human  
129 Transmissible Spongiform Encephalopathies (TSEs) or prion diseases. The mortality rate of TSEs  
130 ranges approximately from 1.5 to 2 persons per million population per year. TSEs can occur  
131 sporadically (sporadic CJD (sCJD)), variably proteinase sensitive prionopathy and sporadic fatal



132 insomnia), be associated with mutations of the prion protein gene (genetic TSEs (gTSE)), or result  
133 from medical exposure to infectious material (iatrogenic CJD (iCJD)). In 1996, a variant form of CJD  
134 (vCJD) was identified<sup>1</sup>. There is strong evidence that vCJD is caused by the agent responsible for  
135 bovine spongiform encephalopathy (BSE) in cattle<sup>2,3,4</sup>. The most likely hypothesis is that vCJD has  
136 occurred through exposure to BSE contaminated food.

137 Human TSEs, including in particular vCJD, were addressed in expert meetings/workshops at the EMA in  
138 January 1998, January 1999, December 1999, May 2000, and December 2000<sup>5c, 5d, 5e</sup>. A CPMP Position  
139 Statement on variant CJD and plasma-derived medicinal products was issued in February 1998<sup>5b</sup> and  
140 the outcome of the subsequent meetings was published on the EMA website. An EMA Expert Workshop  
141 on Human TSEs and Medicinal Products was held on 19-21 June 2002. This provided the scientific basis  
142 for a new CPMP Position Statement issued in 2003<sup>5b</sup>. A further EMA Expert Workshop was held in  
143 January 2004 to review the current state of knowledge of vCJD, in the light of a report of a possible  
144 human transmission by blood transfusion<sup>6</sup>. In addition, the Workshop discussed the CPMP Discussion  
145 document on the investigation of manufacturing processes with respect to vCJD<sup>5a</sup>. In October 2005, a  
146 follow-up workshop was held to discuss the number of vCJD cases reported in France and other  
147 European countries and the potential effect of additional donor exclusion measures. Urine-derived  
148 medicinal products were specifically discussed at an EMA expert workshop in July 2007<sup>5g</sup> after  
149 publication of experiments indicating transmission of infection via urine using a hamster model. A  
150 revised version of the CPMP position statement was published in 2011<sup>5h</sup>.

151 Blood and blood components for transfusion are outside the scope of this Position Statement.  
152 Recommendations on the suitability of blood and plasma donors and the screening of donated blood in  
153 the European Community were described in Council Recommendation 98/463/EC<sup>7c</sup>. European  
154 legislation on human blood and blood components entered into force on 8 February 2003<sup>7a</sup>. Under this  
155 legislation, a Commission Directive on certain technical requirements for blood and blood components,  
156 including eligibility criteria for donors, entered into force in April 2004<sup>7b</sup>.

157 In addition, Council of Europe Recommendation No. R (95) 16 contains a technical appendix on the  
158 use, preparation and quality assurance of blood components and details the current requirements for  
159 donors<sup>8</sup>.

160 In December 2003, following the announcement of a possible case of vCJD transmission by blood  
161 transfusion, Commissioner Byrne made a statement highlighting EU activities in the area of vCJD and  
162 announcing a meeting of the Working Group of the Blood Regulatory Committee to consider the latest  
163 information available from the UK<sup>7d</sup>. The meeting took place in January 2004 and a summary  
164 statement was produced<sup>7e</sup>.

165 The Scientific Steering Committee (SSC), the Scientific Committee on Medicinal Products and Medical  
166 Devices (SCMPMD) and the Scientific Committee on Emerging and Newly Identified Health Risks  
167 (SCENIHR) of the European Commission have published a number of opinions relating to TSEs, which  
168 are of relevance to blood and blood components for transfusion, as well as to plasma-derived medicinal  
169 products<sup>9</sup>. WHO Guidelines on TSEs are also of relevance to both blood components for transfusion and  
170 plasma-derived medicinal products as well as urine-derived medicinal products<sup>10</sup>. The Council of  
171 Europe has made recommendations for blood and blood components for transfusion<sup>11</sup>.

172 The purpose of this revision is to update the position statement according to the recent scientific  
173 developments since the last revision in 2011. This included developments in detection techniques,  
174 epidemiological studies/findings, studies on the tissue distribution of (v)CJD agent, and a study  
175 indicating blood from some patients with sCJD might be infectious.

## 176 **2. Human TSEs current status**

### 177 **2.1. Sporadic, genetic and iatrogenic forms of human TSEs**

178 There is no firm evidence that sporadic, genetic or iatrogenic forms of human TSEs have been  
179 transmitted from person to person through exposure to plasma products or urinary derived medicinal  
180 products. Systematic surveillance for CJD of all types has been undertaken in a number of countries,  
181 including a collaborative study in the EU since 1993,<sup>12,13</sup> and no case of sporadic, genetic or iatrogenic  
182 CJD has been causally linked to prior treatment with plasma products. Two plasma product recipients  
183 in the UK have been diagnosed with sporadic CJD<sup>14</sup>. Both were aged 64 years and had been exposed  
184 to UK sourced plasma products, one for the treatment of von Willebrand's disease and the other  
185 Haemophilia B. Both patients had received, in addition to plasma products, multiple blood transfusions,  
186 but a partial look-back study performed for one patient has not identified a donor with either sCJD or  
187 vCJD. A causal link between the treatment with plasma products and the development of sCJD has not  
188 yet been established and there is a possibility that both cases may reflect a chance event in the  
189 context of systematic surveillance of CJD in large populations<sup>14</sup>.

190 Cases of sporadic CJD with a history of drug treatment for infertility have not been identified but there  
191 is uncertainty about the validity of this observation (see the report of the 2007 EMA expert meeting for  
192 further details)<sup>59</sup>. The strength of epidemiological evidence excluding transmission by urinary derived  
193 medicinal products is less secure than in other forms of human prion disease.

194 Variably proteinase sensitive prionopathy (VPSPr) is an idiopathic disorder with patients having no  
195 known risk factors for acquired or genetic prion disease. Recent laboratory studies have indicated  
196 limited transmissibility to transgenic mice, with transmission characteristics distinct from sporadic  
197 CJD<sup>15, 16</sup>.

### 198 **2.2. Variant CJD**

199 The official UK figures for vCJD at the end of May 2016 were a total of 178 definite or probable vCJD  
200 cases<sup>17</sup>. (One case diagnosed in Hong Kong was classified as a UK case and is included in the UK  
201 figures.) Outside of the UK, there have been 27 cases in France<sup>18</sup>, 5 in Spain, 4 in the Republic of  
202 Ireland and the USA, 3 in the Netherlands and Italy, 2 in Portugal and Canada and single cases in  
203 Saudi Arabia, Japan and Taiwan. Some of these cases, 2 of the Irish cases, 2 of the US cases, 1 French  
204 case, 1 Canadian case and the Taiwanese case had spent more than 6 months in the UK during the  
205 period 1980-1996 and were probably infected while in the UK<sup>19</sup>. The third and fourth US cases and the  
206 second Canadian case have been reported as most likely infected when living outside the USA. The  
207 possibility of cases occurring in other countries cannot be excluded.

208 Two cases of vCJD identified in Spain occurred in the same family. No family links have been reported  
209 in any other vCJD cases to date.

210 All definite and probable cases genotyped had been Met-Met homozygotes at codon 129 of the prion  
211 protein (PrP) gene<sup>20</sup>. In 2016, a definite case of variant CJD was reported in the UK with a  
212 heterozygous codon 129 genotype, raising the possibility of a further outbreak of cases in this genetic  
213 background<sup>21</sup>.

214 Analysis of the figures indicates that vCJD incidence in the UK and internationally is in decline.  
215 However, single cases of vCJD have been identified in the UK and Italy<sup>22</sup> in 2016 and there may be a  
216 long tail or more than one peak to the epidemic.

217 A UK study screening specimens from surgically removed appendices and tonsils for accumulation of  
218 disease related prion protein in the lymphoreticular system, has been carried out in order to try and  
219 obtain some estimation of the number of people that might be incubating vCJD in the UK<sup>23</sup>. Three  
220 positive appendix specimens have been found as a result of the screening of 12,674 appendix and  
221 tonsil specimens. However, the pattern of lymphoreticular accumulation in two of these samples was  
222 dissimilar from that seen in known cases of vCJD, raising the possibility that they may be false  
223 positives. With respect to this possibility, the authors comment that although it is uncertain whether  
224 immunohistochemical accumulation of disease-related prion protein in the lymphoreticular system is  
225 specific for vCJD, it has not been described in any other disease, including other forms of human prion  
226 disease or a range of inflammatory and infective conditions. Subsequent genetic analysis of residual  
227 tissue samples from these 2 cases found that both were valine homozygotes at codon 129 in the prion  
228 protein gene<sup>24</sup>. This finding might account for the immunohistochemical features in these cases; none  
229 of the patients who have developed vCJD and have undergone a comparable genetic analysis have  
230 been valine homozygotes at codon 129 in the prion protein gene.

231 Statistical analysis on this finding of 3 positive specimens gives the following estimations of numbers  
232 who may be incubating vCJD in the UK:

233 237 infections per million population (95% confidence interval (CI): 49-692 per million)

234 These estimations are higher than predictions from modelling of the clinical data (upper 95%  
235 confidence interval of 540 future cases)<sup>25</sup>. It is not known whether those incubating vCJD will  
236 eventually develop clinical disease. However, estimates of numbers possibly incubating are important  
237 with respect to any potential for secondary transmission (e.g. by blood donation, surgical instruments)  
238 while individuals are in the incubation phase. It should be noted that plasma-derived medicinal  
239 products have not been manufactured from donations collected in the UK since 1998.

240 A larger study of an archive of tonsil tissue from 63,007 people of all ages removed during routine  
241 tonsillectomies has been published<sup>26</sup>. In this study, 12,753 samples were from the 1961- 1985 birth  
242 cohort in which most cases of vCJD have arisen and 19,808 were from the 1986-1995 birth cohort,  
243 that may also have been orally exposed to bovine spongiform encephalopathy. None of the samples  
244 were unequivocally reactive to two enzyme immunoassays and none of the initial reactive samples  
245 were positive for disease-related PrP by immunohistochemistry or immunoblotting. The estimated 95%  
246 confidence interval for the prevalence of disease-related PrP in the 1961-1995 birth cohort was 0-113  
247 per million and in the 1961-1985 birth cohort, 0-289 per million. These estimates are lower than the  
248 previous study of appendix tissue, but are still consistent with that study. To confirm the reliability of  
249 the results from the 1961-85 birth cohort, 10,075 of these samples were investigated further by  
250 immunohistochemistry on paraffin-embedded tonsil tissues using two anti-PrP monoclonal antibodies<sup>27</sup>.  
251 One specimen showed a single positive follicle with both antibodies on 2 slides from adjacent sections,  
252 although the earlier enzyme immunoassays and immunoblotting studies on the frozen tissue samples  
253 from this case were negative<sup>26, 27</sup>. If this case is now accepted as positive for abnormal PrP (since the  
254 findings were similar to those of the three positive cases in the earlier study of Hilton et al in 2004<sup>23</sup>),  
255 it gives a prevalence of disease-related PrP in the UK population of 109 per million, with a 95%  
256 confidence interval of 3-608 per million, which is not statistically significantly different (exact  $p = 0.63$ )  
257 from the population prevalence based on the finding of 3 positives in the Hilton et al study<sup>23, 27</sup>. If the  
258 case is not accepted as a positive, this gives a prevalence of 0 out of 9160, with a 95% confidence  
259 interval of 0-403 per million for the 1961-85 cohort, which is also not significantly different (exact  $p =$   
260  $0.25$ ) from the findings of the Hilton et al study<sup>23</sup>. A more recent study from 2013 included 32,441  
261 appendix samples and 16 were positive leading to an estimated prevalence in the UK population of 492



262 cases per million, with wide confidence intervals. All three PRNP codon 129 genotypes were identified  
263 among the 16 positive samples with a relative excess of the VV genotype<sup>28</sup>.

264 The results of further UK prevalence studies of appendix tissues derived from individuals either before  
265 the BSE epidemic or after the introduction of further measures to restrict BSE in the food chain have  
266 recently been published<sup>29</sup>. Positives were found in both groups and the report concluded: "the  
267 Appendix-III survey data have not produced a clear answer to the question of whether abnormal  
268 prions detected by immunohistochemistry in the British population is limited to those exposed to the  
269 BSE epizootic, and various interpretations are possible<sup>29</sup>.

### 270 **3. Human tissue distribution of infectivity/abnormal prion** 271 **protein.**

272 Tissue distribution has been investigated by detection of the abnormal prion protein (PrP<sup>TSE</sup>) or by  
273 infectivity assays. Detection of PrP<sup>TSE</sup> in tissues has often been associated with infectivity, however it  
274 should be noted that infectivity can be present without detection of PrP<sup>TSE</sup>,<sup>30</sup> or PrP<sup>TSE</sup> be present in the  
275 absence of infectivity<sup>31</sup> and that the relation between the amount of PrP<sup>TSE</sup> and infectivity is strain  
276 dependent<sup>32</sup>. The reason for this finding is not known but may be related to limitations of assay  
277 methods for PrP<sup>TSE</sup> or different ratios between protease-resistant and protease-sensitive PrP<sup>TSE</sup>  
278 isoforms<sup>33,34</sup>. It is thus recommended that any study on tissue or fluid distribution of the abnormal  
279 prion protein be confirmed with an infectivity assay.

280 A wider distribution and higher level of PrP<sup>TSE</sup> in human peripheral tissues, including the  
281 lymphoreticular system, has been found in vCJD compared with sporadic CJD<sup>35, 36, 37</sup>. The magnitude of  
282 PrP<sup>TSE</sup> may vary however, as a recent case of vCJD reported extremely low levels of PrP<sup>TSE</sup> in  
283 lymphoreticular tissues<sup>38</sup> and recent data showed equal amounts of PrP<sup>TSE</sup> in vCJD and sporadic CJD<sup>39</sup>.  
284 Limited data from infectivity assays of vCJD tissues are consistent with the PrP<sup>TSE</sup> findings<sup>40</sup>. In clinical  
285 vCJD cases, high titres of infectivity are found in the brain and spinal cord and lower levels in spleen  
286 and tonsil<sup>40, 41</sup>. Infectious vCJD infectivity was detected in spleen but not in the brain from an individual  
287 with the methionine-valine (MV) genotype<sup>42</sup>. While PrP<sup>TSE</sup> and infectivity are occasionally found in the  
288 spleen of sporadic CJD, the levels of PrP<sup>TSE</sup> are lower than in vCJD. PrP<sup>TSE</sup> accumulations have been  
289 observed in muscles of some patients with both sporadic and variant CJD<sup>43</sup>.

290 One study reported that the distribution of PrP<sup>TSE</sup> in iCJD is more similar to sCJD than vCJD<sup>36</sup>. Data are  
291 lacking for gCJD.

### 292 **4. Infectivity in blood and transmissibility via blood**

#### 293 **4.1. Animal blood**

294 In early 2000, most of the knowledge relating to the presence of prion infectivity in blood relied on  
295 information from rodent prion disease models. In these experimental systems, prion infectivity titres  
296 were reported to vary between 1 and 10 ID<sub>50</sub>/mL of blood during the asymptomatic phase and up to  
297 100 ID<sub>50</sub>/mL during the clinical phase of the disease<sup>44, 45</sup>. In these bioassays, infectious prion titres  
298 were measured by bioassay performing intracerebral inoculation of blood, or blood fractions from the  
299 same animal species to indicator animals, (i.e. autologous combinations of inocula and animal  
300 bioassay). The observed infectious prion titres were equivalent to the level of infectivity found in 10<sup>-6</sup> -  
301 10<sup>-8</sup> g of brain tissue from animals at the terminal stage of prion disease. It was found that  
302 approximately 40% of the prion infectivity was associated with the buffy coat fraction, the remainder



303 was found principally in plasma<sup>46, 47</sup>. Importantly, buffy coat-associated prion infectivity was reportedly  
304 washed off these cells by rinsing with PBS.<sup>48</sup> Platelets were shown to have little, if any, prion  
305 infectivity<sup>49</sup>.

306 Subsequent experiments in other animal species, whereby donor blood material was assessed by  
307 bioassay in a host via intracerebral inoculation, have investigated the distribution of prion infectivity in  
308 various blood fractions. Infectivity has also been detected in buffy coat of a prosimian microcebe<sup>50</sup> and  
309 in whole blood of a macaque experimentally infected with a macaque-adapted BSE strain<sup>51</sup> and in red  
310 blood cells of two macaques experimentally infected with a macaque-adapted vCJD strain<sup>51</sup>. In sheep,  
311 naturally or experimentally infected with scrapie, infectious prion titres in whole blood were similar to  
312 those observed in rodents (<35 ID<sub>50</sub>/mL) when measured by bioassay in reporter ovine PrP transgenic  
313 mice<sup>52</sup>. Prion infectivity was detected in plasma from scrapie-infected sheep, but at a lower proportion  
314 to that found in the blood of prion-diseased mice and hamster models<sup>53</sup>. Moreover, a substantial level  
315 of prion infectivity was detected in sheep platelets and infectivity associated with leukocytes was not  
316 reduced by washing of these cells<sup>52</sup>. Similar observations were reported in deer naturally infected with  
317 chronic wasting disease<sup>54</sup>.

318 The intracerebral inoculation of prions is unlikely to recapitulate the cellular and molecular events that  
319 occur as a consequence of prion infection by blood transfusion, a process that involves the  
320 administration of large numbers of viable cells and/or a large volume of material intravenously injected  
321 into the recipient.

322 The relative similarity in size between sheep and humans allows the transfusion of ruminant blood  
323 volumes that are relevant to human medicine. In addition, the pathogenesis of vCJD mirrors features  
324 similar to natural classical scrapie in sheep, for example the presence of prions in peripheral lymphoid  
325 tissue of affected individuals. Consequently, sheep prion disease models were considered to be  
326 relevant models for the assessment of the risks associated with vCJD blood-borne transmission<sup>55</sup>.

327 In early 2000, transfusion of whole blood collected from asymptomatic sheep infected with either  
328 natural scrapie or experimental BSE resulted in prion transmission to recipient sheep<sup>56, 57</sup>.

329 Using the sheep transfusion model, it was also confirmed that RBCs, plasma, platelets and buffy coat  
330 prepared by similar protocols to those used in transfusion medicine can transmit prion disease<sup>58, 59</sup>. In  
331 two different sheep scrapie models, the transfusion of 200 mL of whole blood collected during the early  
332 preclinical phase of the condition (3 months post infection) was able to transmit the disease with 100%  
333 efficacy<sup>52, 58</sup>. However, in two other sheep prion disease studies, the efficacy to transmission after  
334 transfusion of ca. 400 mL of whole blood at a late stage of incubation of the disease was limited to  
335 19%<sup>57</sup> or 40%<sup>59</sup> respectively<sup>57</sup>. Features of the different sheep prion disease models, such as age of  
336 animals used, PrP genotype of the animals and/or the prion strain used for inoculation could contribute  
337 to an explanation for the discrepancies between the results of these different models. However, these  
338 sheep blood transfusion studies collectively suggest that in a proportion of prion-infected blood donors,  
339 the level of prionemia may be insufficient to allow prion disease transmission by blood transfusion<sup>60</sup>.

340 Transfusion experiments carried out in a sheep scrapie model demonstrated that the transfusion of 200  
341 µL of prion-infected whole blood has an apparent 100% efficacy for disease transmission and that  
342 100µL blood transfusion is still sufficient to transmit the disease in a proportion of the recipients<sup>53</sup>.  
343 These experiments also indicated that, despite their apparent low infectious titre, the intravenous  
344 administration of white blood cells (WBC) resulted in efficient disease transmission. The intravenous  
345 administration of 10<sup>5</sup> WBCs were sufficient to cause scrapie in recipient sheep. Cell-sorted CD45R+  
346 (predominantly B lymphocytes), CD4+/CD8+ (T lymphocytes) and CD14+ (monocytes/macrophages)  
347 blood cell sub-populations were all shown to contain prion infectivity by bioassays in ovine PrP

348 transgenic mice<sup>51</sup>. However, while the intravenous administration of 10<sup>6</sup> CD45+ or CD4/8+ living cells  
349 were able to transmit the disease, similar numbers of CD14+ failed to infect any of their recipients.  
350 These indicated that blood cell populations display different abilities to transmit TSE by the transfusion  
351 route.

352 PrP<sup>TSE</sup> has been detected in blood components of TSE-infected animals by different techniques. In TSE-  
353 infected rodents, PrP<sup>TSE</sup> positivity has been reported in buffy coat<sup>62</sup> and plasma exosomes<sup>63</sup> by Protein  
354 Misfolding Cyclic Amplification (PMCA), whole blood by Real-Time Quaking induced Conversion Assay  
355 (RT-QuIC)<sup>64</sup>, and by steel-binding assay<sup>65</sup> and in plasma exosomes by standard Western Blot (WB)  
356 procedures.<sup>66</sup> Abnormal PrP conformers can be detected throughout the whole incubation period of the  
357 disease<sup>65</sup>.

358 In pre-clinical and clinical scrapie-infected or BSE infected sheep, PrP<sup>TSE</sup> positivity has been reported in  
359 platelets and WBC by PMCA or infectivity assay<sup>52,67,68</sup> or surface-FIDA (fluorescence intensity  
360 distribution analysis)<sup>68</sup>. In chronic wasting disease (CWD)-infected deer, whole blood resulted PrP<sup>TSE</sup>  
361 positive by RT-QuIC in both animals in both, pre-clinical and clinical phases of disease<sup>63</sup>. Plasma, buffy  
362 coat and WBC tested PrP<sup>TSE</sup> positive by PMCA in vCJD-infected macaques during the earliest pre-clinical  
363 and clinical phases of disease<sup>66,70, 71</sup>.

#### 364 **4.2. Human blood**

365 The tracing of recipients of blood transfusion from UK donors who have subsequently developed vCJD  
366 (the Transfusion Medicine Epidemiology Medicine Review, TMER study) has revealed four instances of  
367 secondary transmission<sup>72</sup>. These individuals had received transfusion of non-leucodepleted red cells  
368 from donors who were clinically healthy at the time of donation but subsequently (17–40 months later)  
369 developed variant CJD. Three of the four patients developed disease after incubation periods ranging  
370 from 6.5 to 8.5 years; the fourth died of an illness unrelated to prion disease 5 years after transfusion.  
371 This asymptomatic prion-infected patient was heterozygous (methionine/valine) at codon 129 of  
372 the *PRNP* gene. However the spleen and lymph nodes tested positive<sup>73</sup> and the prion agent was  
373 experimentally transmitted from brain and spleen to humanised transgenic mice<sup>74</sup>. Taken together,  
374 these instances are strong evidence that vCJD is transmissible through blood transfusion.

375 In 2010, another presumed case of asymptomatic vCJD infection was identified in an elderly  
376 haemophilia patient who was heterozygous at codon 129 in the prion protein gene<sup>75</sup>. The patient, who  
377 died of unrelated pathology, had received large quantities of UK-sourced fractionated plasma products  
378 (i.e. FVIII), including some units derived from plasma pools which contained plasma from a donor who  
379 later developed variant CJD. This patient was identified through an intensive search for PrP<sup>TSE</sup> positivity  
380 in a range of post-mortem tissues, although only 1 of 24 samples taken from the spleen tested  
381 positive. Whether someone with this limited distribution of PrP<sup>TSE</sup> would be infectious is unknown, but  
382 from a public health perspective, this patient represents a warning that some plasma-derived products  
383 might contain residual prion infectivity.

384 The surveillance described above emphasises the importance of the TMER study for identifying the risk  
385 of blood transfusion in transmitting vCJD. Moreover, national databases of blood donors and the  
386 maintenance of traceability from donor to recipient and vice versa are essential to establish whether a  
387 vCJD case has been a blood donor (UK experience has shown that questioning of family members is  
388 unreliable for establishing whether a patient has been a blood donor). Traceability is a specific  
389 requirement in Article 14 of Directive 2002/98/EC.

390 In a conventional mouse model (RIII mice), infectivity was not detected in the blood of two vCJD cases  
391 but the bioassay had limited sensitivity to detect infectivity in peripheral tissues such as tonsil or  
392 spleen<sup>40</sup>. Bioassays carried out in PrP transgenic mice using blood harvested post mortem from a vCJD-  
393 affected patient have shown the presence of prion infectivity in red blood cells, plasma and white blood  
394 cells<sup>74</sup>. The blood fractions used in these assays had been prepared in 2000 using laboratory-scale  
395 haematological protocols but did not include leukoreduction. The infectious titre of whole blood in the  
396 bioassayed vCJD sample was estimated to be approximately 4.45 ID<sub>50</sub>/mL, which is 10<sup>-6</sup> - 10<sup>-7</sup> lower  
397 than that found in one gram of brain from a vCJD-affected patient at terminal stage of disease.  
398 Importantly, the leukocyte-associated prion infectivity of the vCJD blood sample could not be reduced  
399 by rinsing of the cells, similar to that found in ruminant animal models. These data support the view  
400 that prion infectivity levels in the blood of vCJD patients and different animal prion disease models are  
401 similar. They also demonstrated that interspecies variations exist with regards to distribution of  
402 infectivity in different blood fractions.

403 Look-back studies in the UK<sup>77</sup> and USA<sup>78</sup> have not revealed any possible case of sporadic CJD linked to  
404 blood transfusion. However, current data are too scant to unequivocally exclude the possibility that  
405 such an event could occur in a small number of cases with a long (10 or more years) incubation period.

406 A review of transmission studies to detect infectivity in the blood of humans with sporadic and  
407 iatrogenic CJD shows that experimental transmissions in animal models have occasionally been  
408 reported in some studies<sup>79-83</sup> but not in others.<sup>84</sup> It is possible that PrP<sup>TSE</sup> is present at low levels in the  
409 blood of clinically affected cases of sCJD. Recently, intracerebral inoculation of plasma from two of four  
410 sporadic CJD patients transmitted disease into human PrP transgenic mice. The relative infectivity  
411 between brain and plasma was the same in sCJD and vCJD<sup>76</sup>. Data are lacking for gCJD and iCJD.

412 PrP<sup>TSE</sup> was detected in WBC of a single vCJD patient, in buffy coat of 2 out of 3 vCJD patients by  
413 PMCA<sup>67</sup> and in the blood of 15 out of 21 vCJD cases by steel binding assay<sup>85</sup>.

414 For the purpose of risk assessments, it is recommended that, as a worst case assumption, a relative  
415 efficiency of the intravenous and intracerebral routes of 1:1 should be used.<sup>86</sup> This is because the  
416 accumulated information now available from animal studies indicates that the intravenous route can be  
417 an efficient route of transmission and in certain cases can give a transmission rate and/or an  
418 incubation period similar to the intracerebral route (see also 4.1).

## 419 **5. Detection techniques**

420 A donor screening test could provide an improved level of safety. The development of blood tests for  
421 vCJD remains a strategic priority but has suffered from declining efforts from an assumption that the  
422 technical challenges are insurmountable, an assumption that has seen commercial bodies abandoning  
423 test development<sup>87</sup>.

424 As unique biological agents mammalian prions provide many research challenges. Not least is the  
425 ability to detect and quantify their presence in tissue and fluid samples. The severity of pathology  
426 associated with clinical prion disease suggests markers for infection and disease progression other than  
427 abnormal PrP may exist. Numerous studies by groups worldwide<sup>88-94</sup> have applied 'omics' approaches  
428 to discovery of alternative markers. Several differential changes between baseline and disease states  
429 have been demonstrated but they lack the specificity required for use in screening or diagnostic tests<sup>8</sup>.  
430 In contrast the deposition of PrP<sup>TSE</sup> is the archetypal marker of prion disease. Whilst moderately  
431 abundant in the tissues of the central nervous system and lymphoreticular tissue in cases of vCJD, the  
432 concentration of infectivity, and by inference PrP<sup>TSE</sup>, is very low in blood and cerebrospinal fluid (CSF).



433 This situation is further complicated by the large background excess of normal non-pathogenic cellular  
434 protein PrP<sup>C</sup> associated with the cellular compartment of blood.

435 A conceptually obvious approach to overcome the problems of abnormal PrP detection is to exploit the  
436 innate propensity of amyloid to self-propagate. This approach has been developed in a variety of  
437 formats of which two: QuIC<sup>95</sup> and PMCA<sup>96</sup> have seen widespread adoption and development for  
438 research. The adoption of QuIC for the diagnosis of sporadic CJD using CSF samples has been  
439 successful with excellent although not perfect performance characteristics<sup>97</sup>. However, adaptation of  
440 this methodology to the testing of blood samples has yet to be convincingly demonstrated. PMCA has  
441 been shown to be capable of detecting vCJD infection in blood<sup>67</sup> and urine<sup>98</sup>. However, the specificity of  
442 such an assay is generally considered to be a frailty of this approach. Two recent studies using PCMA  
443 showed 100% sensitivity at identification of blood samples from 14<sup>99</sup> or 18<sup>100</sup> clinical vCJD cases and  
444 indicated specificities in the range as required in the EU Common technical specification (CTS)<sup>101</sup>.  
445 However, full validation according to the CTS has not yet been performed.

446 As an alternative to amplification strategies, enrichment by capture using stainless steel beads has  
447 allowed the direct immunoassay of captured material, detecting a signal in blood in 71% (15 out of 21)  
448 of vCJD patients<sup>85</sup> whilst being highly specific<sup>102</sup>.

449 It is clear that there several methods in research and development that offer possibilities for routine  
450 screening and confirmatory assays but they have not yet completely demonstrated the current  
451 requirements of sensitivity and specificity as defined in the Common Technical Specifications.<sup>101</sup>  
452 Comparison and validation of potential screening tests is considerably confounded by the paucity of  
453 blood samples from confirmed cases of clinical prion disease and very limited samples available from  
454 asymptomatic individuals who later developed vCJD.

## 455 **6. Leucoreduction and specific prion affinity filters**

456 Leucodepletion was introduced in the UK in 1999 as a precautionary measure in transfusion medicine  
457 to reduce the risk of iatrogenic transmissions of vCJD. The rationale was based upon evidence to  
458 suggest the majority of infectivity in whole blood is associated with 'buffy coat' fractions or  
459 mononuclear cells.

460 Despite widespread exposure to potentially contaminated blood transfusions in the UK, Europe and the  
461 wider world, confirmed cases of vCJD resulting from exposure to contaminated blood or blood products  
462 are small<sup>75, 103, 104</sup>. This may be partly attributed to the rapid introduction of leucodepletion.

463 In addition to the potential protection afforded against vCJD transmission, leucodepletion has other  
464 benefits in transfusion medicine including reduced risk of HLA alloimmunisation with the potential for  
465 refractoriness to platelet transfusion, reduction in specific viral transmission risk, the disappearance of  
466 transfusion-related graft versus host disease and a significant decrease in cases of post-transfusion  
467 purpura<sup>105</sup>.

468 Experience from animal models indicates that leucodepletion is highly effective for prion safety of blood  
469 transfusion. Taken together with the additional benefit of improved red blood cell and platelet quality it  
470 is clear that leucodepletion is advantageous and is likely to remain in place irrespective of prion  
471 transmission risk assessments.

472 The Scientific Committee on Medicinal Products and Medical Devices (SCMPMD) opinion on  
473 leucoreduction<sup>9a, 9b</sup> for blood and blood components for transfusion stated that it might be a



474 precautionary step to remove white cells as completely as possible. For plasma for fractionation the  
475 opinion stated the following:

476 'Taken together, there is no compelling scientific evidence to date for the introduction of leucoreduction  
477 of plasma for fractionation, or other methods aiming at removal of cells and debris, as a precaution  
478 against vCJD transmission. The question should be further explored by suitable experiments.'

479 Results reported at the 2002 EMEA Workshop, suggested that leucodepletion does not cause  
480 fragmentation of cells and lysis. Results of a comprehensive study involving a number of different  
481 filters and procedures indicate that leucodepletion is not detrimental in terms of the generation of  
482 microvesicles or the release of prion proteins<sup>106</sup>.

483 Specific affinity ligands that bind prion proteins have been evaluated for their ability to further reduce  
484 TSE infectivity present in blood and plasma. Exogenous spiking experiments have suggested prion-  
485 specific filters could be effective. However, such studies do not provide a good model of infectivity  
486 distribution in blood and endogenous validation experiments have indicated the efficiency of prion  
487 removal is not very effective with an overall logarithmic reduction value of only 1.22 from infectivity  
488 assay in a hamster model<sup>107</sup>.

489 In October 2009, the UK Advisory Committee on the Safety of Blood, Tissues and Organs (SaBTO)  
490 stated that there was sufficient evidence that a specific affinity ligand filter reduces infectivity and  
491 recommended the use of prion filtration of red cell components administered to children born since 1  
492 January 1996. This recommendation was subject to the satisfactory completion of the PRISM clinical  
493 trial to evaluate the safety of prion filtered red blood cells<sup>108</sup>.

494 Despite the fact that PRISM has indicated that the use of commercially available prion filters was not  
495 detrimental to the quality or safety of filtered red blood cells, the use of prion reduction filters has not  
496 been recommended. This decision has been based upon the need for independent studies to replicate  
497 the findings of these studies since the studies involved the filter manufacturers.

498 Two such studies were commissioned and finally published in 2015. One, using a hamster model of  
499 prion disease concluded that the majority of infectivity was removed using leucodepletion alone, with  
500 filtration using the CE marked prion filter P-Capt (MacoPharma, France) achieving a further reduction  
501 in titre of around only 0.2 ID/ml.<sup>109</sup> The study was compromised by the low dynamic range afforded by  
502 the input material, however, residual infectivity was still present following combined leucodepletion and  
503 prion filtration and the low concentration was not statistically different from the residual levels  
504 following leucodepletion alone. The second study involved transfusion from scrapie-infected sheep and  
505 recipients received either leucodepleted blood or sequentially leucodepleted and P-Capt prion filtered  
506 blood<sup>110</sup>. This study also concluded that there was no significant difference in residual titre following  
507 only leucodepletion or leucodepletion and prion filtration. However, this study was also flawed in that  
508 all transfused materials were leucodepleted and the genotypes of recipient sheep were not disclosed so  
509 the possibility of resistant genotypes being transfused cannot be excluded. As a result, despite the  
510 large number of sheep used in the study, only two recipient animals were considered transfusion  
511 positive; one having received leucodepleted blood and the other receiving blood following combined  
512 leucodepletion and prion filtration. In conclusion, both studies failed to demonstrate a clear effect of  
513 the prion affinity filters.

514 The prion binding capacity of another affinity ligand chromatography step has been investigated in the  
515 processing of a plasma medicinal product using hamster brain derived spiking material.<sup>111, 112</sup> These  
516 data require further evaluation before conclusions can be drawn on possible efficacy.

## 517 **7. Manufacturing processes for plasma-derived medicinal** 518 **products**

519 Despite the fact that there is no firm evidence of transmission of CJD through plasma-derived  
520 medicinal products, infectivity has been detected in the plasma of both vCJD and sCJD affected  
521 patients<sup>76</sup>.

522 Taking account of the available data concerning blood infectivity, it is of utmost importance to  
523 investigate the capacities of the manufacturing process (fractionation) to eliminate/inactivate the  
524 infectious material potentially present in the plasma pool used as the starting material for preparation  
525 of plasma-derived products.

526 Initial results from animal studies, using blood from rodents, indicated that the fractionation process  
527 contributes to the decrease of infectivity in some fractionated products<sup>44, 46</sup>.

528 However, information reported at the EMA Workshops in 2002 and 2004 suggested that endogenous,  
529 rodent blood-associated infectivity might persist through the fractionation process to a greater extent  
530 than would be expected from spiking studies using brain-derived prion preparations, possibly because  
531 of the differing physical and biochemical properties of the associated infectious particles.

532 A significant number of studies aimed at following the partition/removal of PrP<sup>TSE</sup> and/or infectivity  
533 during plasma fractionation process have been carried out using such spiking approaches<sup>113, 114</sup>.

534 The vast majority used rodent-adapted TSE agent (263K hamster strain) brain homogenate and  
535 microsomal brain fractions as a spike. They relied on direct PrP<sup>TSE</sup> immunodetection tools (western blot  
536 or conformation dependent immunoassay) to demonstrate a drop in the TSE agent content in  
537 processed fractions and on bioassay infectivity measurements to confirm the results. Generally, the  
538 limited sensitivity of these immuno-detection methods made necessary the use of a massive amount of  
539 TSE agent in the spike.

540 These studies established the potential contribution of the various manufacturing steps to the  
541 reduction of TSE agents (including precipitation followed by centrifugation or depth filtration, specific  
542 chromatographic steps and nanofiltration).

543 However since 2004 and the publication of the EMA guideline on *The investigation of manufacturing*  
544 *processes for plasma-derived medicinal products with regards to vCJD risk* (October 2004), the  
545 knowledge of the prion area in general and the endogenous infectivity in blood in particular, have  
546 significantly evolved. Moreover, experimental studies highlighted the fact that prion removal capacity  
547 may directly vary according to the spiking preparation (dispersion and TSE agents strains) particularly  
548 for steps based on retention mechanisms<sup>115</sup>.

549 These new elements raise questions about the final relevance of certain experimental approaches that  
550 were used for characterizing prion removal capacities of plasma manufacturing steps. Consequently  
551 there is still a need to perform research on the best experimental approach for evaluation of the  
552 partitioning or removal capacities of the various fractionation steps used in the preparation of plasma-  
553 derived medicinal products.

554 It is recommended to use various forms of spike preparations in order to obtain an insight into their  
555 influence on prion reduction at the specific investigated step as compared to what has been published  
556 in the literature. In specific cases, it might be worth considering the use of blood from infected animals  
557 as an alternative material for investigation of early plasma processing steps, where feasible and where  
558 the overall prion reduction capacity seems limited or questionable. There is still further need for

559 research to gain better knowledge of the form of infectivity present in blood (or in intermediates from  
560 manufacture) in order to confirm the relevance of the spiking material used in the validation studies.

## 561 **8. Infectivity in urine**

### 562 **8.1. Animal urine**

563 Low levels of infectivity have been detected in urine of scrapie-infected rodents by several research  
564 groups and in the urine of deer with CWD<sup>59</sup>. Accordingly, urine has been reclassified among the  
565 category of "lower-infectivity tissues" by WHO<sup>10c</sup>.

566 Seeger *et al.*<sup>116</sup> have studied transmission via urine using mouse models of chronic inflammation. They  
567 have detected prionuria in scrapie infected mice with coincident chronic lymphocytic nephritis.  
568 Transmission has been shown upon intracerebral inoculation of purified proteins from pooled urine  
569 collected from scrapie sick or presymptomatic mice. In contrast, prionuria was not observed in scrapie  
570 infected mice displaying isolated glomerulonephritis without interstitial lymphofollicular foci or in  
571 scrapie infected wild type mice lacking inflammatory conditions.

572 Gregori *et al.*<sup>117</sup> demonstrated that the disease could be transmitted by intracerebral inoculation of  
573 pooled urine from scrapie-sick hamsters. The infectivity titre of the urine was calculated to be around  
574 3.8 infectious doses/ml. Titration of kidney and urinary bladders from the same animals gave 20,000-  
575 fold greater concentrations. Histologic and immunohistochemical examination of these tissues showed  
576 no indication of inflammation or other pathologic changes, except for occasional deposits of disease-  
577 associated prion protein in kidneys.

578 Prionuria was also detected in CWD of deer. Experiments by Haley *et al.*<sup>118</sup> provided evidence that  
579 concentrated urine from deer at the terminal stage of the disease, that also showed mild to moderate  
580 nephritis histopathologically, was infectious when inoculated into transgenic mice expressing the cervid  
581 PrP gene. In addition, the urine collected from the CWD sick deer that was used for mouse inoculation,  
582 showed positive results when assayed for PrP<sup>TSE</sup> by serial rounds of PMCA assay. The concentration of  
583 abnormal prion protein was very low as indicated by undetectable PrP<sup>TSE</sup> by traditional assays and  
584 prolonged incubation periods and incomplete TSE attack rates in the transgenic mice.

585 Using the highly sensitive PMCA or RT-QuIC technologies, PrP<sup>TSE</sup> have been detected in urine of scrapie  
586 sick hamsters,<sup>119, 120, 121</sup> cervids with preclinical and clinical CWD<sup>122-125</sup> and sheep with at preclinical  
587 and clinical stages of scrapie disease scrapie<sup>125</sup>. The concentration of PrP<sup>TSE</sup> in urine is, on average, 10-  
588 fold lower than in blood<sup>119</sup>.

### 589 **8.2. Human urine**

590 Epidemiological evidence in the last 25 years, during which urinary-derived medicinal products and  
591 particularly gonadotrophins have been widely used, does not suggest, at present, a risk from sporadic  
592 CJD. Since epidemiological evidence has identified the few cases of iatrogenic transmission of CJD  
593 through the use of pituitary-derived gonadotrophins, it is possible that transmission from urinary-  
594 derived gonadotrophins would have been detected if it had occurred. This is further supported by a  
595 recent study, in which prion infectivity in urine from a sCJD patient was not detected using bioassays in  
596 transgenic mice suggesting that prion infectivity in urine is either not present or was below the  
597 detection limit of 0.38 infectious units/ml<sup>126</sup>.

598 Recently, PrP<sup>TSE</sup> has been detected in the urine of patients with vCJD by using the highly sensitive  
599 PMCA technique<sup>98</sup>, but not in urine of sporadic CJD patients<sup>39, 98</sup>. However the sensitivity of the PMCA



600 detection for sCJD remained unassessed in these studies, raising concern about the significance of  
601 these negative results. More recently, abnormal PrP conformers were also detected in the urine of sCJD  
602 patients using an enrichment technique followed by an immunoassay. In this study, 8 of 20 sCJD cases  
603 tested positive while the analysis of 125 control samples (comprising 91 normal control individuals and  
604 34 neurological disease control individuals), remained negative<sup>127</sup>.

## 605 **9. Recommendations and proposals**

### 606 ***9.1. Sporadic, genetic and iatrogenic CJD and plasma-derived medicinal*** 607 ***products***

608 There is no change in the recommendations for donor selection. There is also no change in the  
609 recommendations for batch recalls. However the importance of the prion-reducing capacity of the  
610 manufacturing process is emphasised.

611 Donor selection criteria include criteria to exclude donors who might be at higher risk of developing  
612 CJD. The following permanent deferral criteria are specified in Commission Directive 2004/33/EC:  
613 Persons who have a family history which places them at risk of developing a TSE, or persons who have  
614 received a corneal or dura mater graft, or who have been treated in the past with medicines made  
615 from human pituitary glands. Precautionary recalls of batches of plasma-derived medicinal products  
616 after post-donation reports of CJD or CJD risk factors in a donor contributed to severe shortages of  
617 certain products<sup>10a</sup>.

618 The perception that plasma products and blood of sporadic CJD patients might contain prion infectivity  
619 has increased because of the recent transmission study with human blood in transgenic mice and the  
620 occurrence of two cases in plasma product recipients. However, cumulative epidemiological evidence  
621 does not support transmission of sporadic, genetic and iatrogenic CJD by blood, blood components or  
622 plasma-derived medicinal products, although the statistical power of these epidemiological studies for  
623 tracing blood-related sCJD cases may not be sufficient to definitively exclude the possibility of blood  
624 transmission in a small number of cases. Therefore, the CHMP recommendation that recall of plasma  
625 derived medicinal products is not justified where a donor is later confirmed as having sporadic genetic  
626 or iatrogenic CJD or risk factors is maintained provided the manufacturer has demonstrated using  
627 appropriate methodology that the process includes steps which will minimize any risk of prion  
628 contamination of the final product.

629 The implementation of appropriate actions in relation to CJD depends on accurate diagnosis in  
630 suspected cases. There is still potential for diagnostic confusion between sporadic and variant CJD,  
631 particularly in younger age groups<sup>128</sup>.

### 632 ***9.2. Variant CJD and plasma-derived medicinal products***

633 There is no change in the recommendations for vCJD. Although the number of cases is in decline in the  
634 UK and France, isolated cases of vCJD are still being reported and there is still uncertainty about the  
635 future number of cases. Variant CJD has a wide distribution of infectivity in tissues outside the central  
636 nervous system.

637 There is strong epidemiological evidence of human to human transmission of vCJD by blood transfusion  
638 (see Section 4.2). In addition, one vCJD infection was detected in a patient with haemophilia treated  
639 with high doses of intermediate purity factor VIII. Estimates of the relative risks of exposure through  
640 diet, surgery, endoscopy, blood transfusion and receipt of UK-sourced plasma products suggest that



641 the most likely route of infection in the patient with haemophilia was receipt of UK plasma products. At  
642 least one batch came from a pool containing a donation from a donor who later developed vCJD.

643 The following measures are aimed at minimising the risk of transmission of the agent by plasma-  
644 derived medicinal products.

### 645 **9.2.1. Exclusion Criteria**

#### 646 **a) Consideration of Country-based exclusions**

647 There is currently no screening test to detect donors who may be incubating the disease or in the early  
648 clinical stages. Therefore, other approaches are considered in order to try and identify donors who may  
649 present a higher risk.

#### 650 ***UK plasma***

651 Residence in the UK is a recognised risk factor for vCJD and has led to the UK deciding no longer to  
652 fractionate from UK plasma.

#### 653 ***Exclusion of donors based on cumulative period of time spent in the UK***

654 Since UK donors are excluded from donating plasma for the manufacture of plasma-derived medicinal  
655 products in the UK, it is consistent to exclude donors who have spent long periods in the UK. This is  
656 supported by the finding of vCJD cases, which have a risk factor of long periods spent in the UK, in  
657 other countries.

658 It is therefore recommended that donors who have spent a cumulative period of 1 year or more in the  
659 UK between the beginning of 1980 and the end of 1996 are excluded from donating blood/plasma for  
660 fractionation. Countries are highly encouraged to choose their national cumulative period limit for  
661 plasma-derived medicinal products according to a nationally calculated benefit/risk balance, which will  
662 take into account the endogenous risk of BSE exposure (and introduction in the food chain) and the  
663 risk of shortages of blood and plasma for the manufacture of medicinal products. The national limit is  
664 recommended to be of cumulative periods in the UK below or equal to 1 year.

665 Countries may still apply a stricter limit than 1 year for exclusion of donors for blood/plasma collected  
666 for fractionation within the country (e.g. 6 months) but will accept plasma-derived medicinal products  
667 from other countries provided that at least the one-year time limit is applied.

668 The rationale for this recommendation is to exclude donors who have the highest individual risk from  
669 stays in the UK and to be consistent with the UK decision to no longer fractionate from UK plasma. This  
670 is further explained in the first version of this Position Statement published in February 2003<sup>5b</sup>.

#### 671 ***French plasma and plasma from other BSE-exposed European countries***

672 France published an analysis of the risk of transmission of vCJD by blood and its derivatives sourced  
673 from French plasma in December 2000<sup>129g</sup>. This concluded that plasma collected in France could  
674 continue to be used for fractionation. The safety margin for plasma-derived medicinal products was  
675 considered to be sufficient. However, introduction of additional steps to further increase the safety  
676 margin of some products was recommended (e.g. nanofiltration of Factor VIII introduced in January  
677 2001). Leucodepletion for plasma for fractionation, as for plasma for transfusion products, was also  
678 recommended in 2001 as a precautionary measure. The subsequent risk-analyses published in 2002,  
679 2003, 2004, 2005, 2007 and 2009 re-confirmed these conclusions and acknowledged that the

680 estimated size of the epidemic had been reduced by more recent modelling, and the risk associated  
681 with collecting blood from vCJD-incubating donors was lower than previously estimated<sup>129</sup>.

682 Based on the limited data on human exposure to BSE-risk materials in other European countries, it is  
683 still difficult to estimate the epidemiological risk in those countries which have a small number of vCJD  
684 cases or have not yet reported any vCJD cases.

685 ***Donors who have spent a cumulative period of time in France and other BSE-exposed***  
686 ***countries***

687 Exclusion of donors who have spent a cumulative period of time in France is not recommended  
688 because of the lower risk associated with time spent in France compared with time spent in the UK  
689 (the risk in France is estimated to be 1/10 of that in the UK)<sup>129b</sup>. Endogenous vCJD cases occurred in  
690 some other countries (see Section 2. Human TSEs current status) placing them close to or lower than  
691 France in terms of incidence and ratio of risk in comparison to UK. Exclusion of donors who have spent  
692 time in other countries having a risk ratio in the same order of magnitude as France is not  
693 recommended.

694 ***Concluding remarks***

695 Country-based exclusions may appear unjustified in the sense that the vast majority of donors who will  
696 be excluded will not develop the disease. There is a lack of spare plasma capacity to make up for  
697 shortfalls if countries that are major producers of plasma-derived medicinal products discontinue the  
698 use of nationally collected plasma for fractionation.

699 **b) Other possible exclusion criteria**

700 Commission Directive 2004/33/EC indicates that further deferral criteria for vCJD may be  
701 recommended as a precautionary measure.

702 Other possible exclusion criteria that could be considered include permanent exclusion of recipients of  
703 blood transfusion in UK.

704 Caution is needed because of the risk of loss of donors and consequent supply problems. Since such  
705 criteria could apply to both blood and blood components, and plasma-derived medicinal products, this  
706 is kept under review within the scope of Directive 2002/98/EC. The Competent Authorities for blood  
707 and blood components expressed the need to have scientific evidence on the safety impact of possible  
708 additional exclusion criteria, as well as to make a national assessment on the expected impact of these  
709 criteria on donation volumes, before implementing additional exclusion criteria.

710 The opinion of May 2006 from the Scientific Committee on Emerging and Newly Identified health Risks  
711 (SCENHIR) stated that it did not consider that additional specific measures were needed to reduce the  
712 risk from vCJD infectivity in blood. When there is a concern for spreading vCJD by blood transfusion,  
713 donor exclusion of blood transfusion recipients is the appropriate measure<sup>9i</sup>.

714 **9.2.2. Leucoreduction and specific prion affinity filters**

715 The benefit of inclusion of leucoreduction to improve the safety of plasma has not been demonstrated.

716 At present it is not appropriate to recommend the introduction of leucoreduction for the safety of  
717 plasma-derived products.

718 Efficacy of introducing recently developed affinity media / filters to blood or plasma has been  
719 investigated. Although they might have some effect in reducing prion loads, clear evidence for their  
720 use in providing protection against transmission is still uncertain.

### 721 **9.2.3. Manufacturing processes for plasma-derived medicinal products**

722 The available data support the reduction of infectivity by steps in the manufacturing process.  
723 Manufacturers are required to estimate the potential of their specific manufacturing processes to  
724 reduce infectivity. This should follow a step-wise approach as described below and illustrated in the  
725 accompanying flow diagram. It is recommended that manufacturers consult the relevant competent  
726 authorities at each of the milestones in this estimation. A decision to add a further manufacturing  
727 step(s) to increase reduction capacity should only be made after careful consideration of all benefit-risk  
728 factors for a certain product.

729 Firstly, manufacturers should compare their own processes to those with published data on reduction  
730 of infectivity in order to estimate the theoretical potential of their specific manufacturing processes to  
731 reduce infectivity. (*Flow diagram, step 1*)

732 Whereas the general information available on manufacturing processes provides useful background  
733 information, the actual effectiveness of a manufacturing process might be dependent on the specific  
734 process conditions. Manufacturers should consider the relevance of the published data to their specific  
735 manufacturing processes and whether the removal capacity can be expected to be comparable.

736 If it cannot be concluded that the removal capacity would be expected to be comparable, it is  
737 recommended that manufacturers undertake product-specific investigational studies on key steps in  
738 their manufacturing processes using biochemical assays. Priority should be given to studies on  
739 products with the lowest potential removal capacity. (*Flow diagram, step 2*)

740 Investigations using biochemical assays may be sufficient if a clear correlation with infectivity data has  
741 already been established for similar processes (e.g. ethanol fractionation). If such a correlation is not  
742 established (e.g. a novel step) and the step is considered critical for removal of infectivity for the  
743 specific product (e.g. it is the only step for removal), the investigations should be confirmed using an  
744 infectivity assay for the critical step(s). (*Flow diagram, step 3*)

745 The above steps will allow manufacturers to estimate the reduction capacity of their manufacturing  
746 processes. (*Flow diagram, step 4*)


747 In cases where the overall reduction capacity is limited, manufacturers should consider the addition of  
748 steps that may increase the removal capacity where this is feasible without compromising the safety,  
749 quality and availability of the existing products. Discussion with the relevant competent authorities is  
750 recommended. (*Flow diagram, step 5*)

751 The outcome of the estimates of the theoretical potential of manufacturing processes to reduce  
752 infectivity and the results of product-specific investigational studies should be reported to the relevant  
753 competent authorities for the medicinal products concerned, as information becomes available.  
754 Applicants submitting new marketing authorisation applications for plasma-derived medicinal products  
755 will be expected to include such information in the application dossier. The outcome of the estimation  
756 of the theoretical potential to reduce infectivity should always be included in the application.

757 In support of these recommendations, CHMP's Biologics Working Party, with the involvement of  
758 external experts, has developed guidance on how to investigate manufacturing processes with regard  
759 to vCJD risk<sup>5a</sup>.

Figure 1: Plasma-Derived Medicinal Products: estimation of potential reduction capacity of specific manufacturing processes

**Important Note:** this flow diagram should be read in conjunction with the preceding text in 9.2.3. It is recommended to consult the relevant competent authorities at the milestones in this estimation. Give priority to studies on products with the lowest potential removal capacity.



760

761 **9.2.4. Recall of batches where information becomes available post-**  
 762 **donation**

763 In view of the lack of adequate information on vCJD, it is prudent to recall batches of plasma-derived  
 764 medicinal products where a donor to a plasma pool subsequently develops vCJD. Recall should also  
 765 include medicinal products containing plasma-derived products as excipients (see also 9.2.5).  
 766 However, in both cases, consequences for essential medicinal products where alternatives are not  
 767 available will need careful consideration by the competent authorities.



768 A case-by-case consideration would be appropriate where plasma-derived products have been used in  
769 the manufacture of other medicinal products. This consideration would include the nature of the  
770 product, the amount used, where it is used in the manufacturing process and the downstream  
771 processing.

772 Look-back to identify the fate of donations should be taken as far as possible. Regulatory authorities,  
773 Official Medicines Control Laboratories, surveillance centres and the supply chain should be informed of  
774 all batches of product and intermediate implicated whether or not supplies of the batch are exhausted.

775 There is no recommendation to recall batches if information becomes available post-donation, which  
776 would have excluded a donor based on his/her stay in the UK (see 9.2.1).

### 777 **9.2.5. Albumin used as an excipient or in manufacturing processes**

778 The available data on the removal of infectivity during the fractionation process used in the  
779 manufacture of albumin indicates that the risk of transmission of infectivity by albumin would be  
780 particularly low. Where a donor to a plasma pool subsequently develops vCJD in the case of albumin  
781 used as an excipient, a recall should be considered. However, a careful case-by-case risk analysis  
782 taking into account the estimated capacity of the process to remove infectivity and the amount of  
783 albumin incorporated in the medicinal product could justify not needing a recall. A single batch of  
784 albumin may be used to produce a number of batches of a medicinal product because of the small  
785 amounts that are typically used as an excipient. As a consequence, a recall could affect complete  
786 stocks of a product and create severe shortages. Therefore, to avoid a negative impact on supply,  
787 companies should consider the origin of plasma and select countries where the probability of having to  
788 recall batches is as limited as possible.

789 Use of substitutes for plasma-derived albumin used as an excipient or in manufacturing processes is  
790 encouraged and should be considered as a long-term approach.

### 791 **9.2.6. Substitution with alternative products**

792 Use of alternative products to plasma-derived medicinal products could be considered, where these are  
793 available. It is felt that this choice should remain with users, taking into account the needs of the  
794 individual patient. It should be noted that plasma-derived products such as albumin may be used in  
795 the manufacture of recombinant products.

### 796 **9.2.7. Optimal Use**

797 Optimal use of plasma-derived medicinal products is encouraged, as this will maximise the benefits of  
798 the products compared with any potential risk.

## 799 **9.3. Urine-derived medicinal products**

800 The recommendations for urine-derived medicinal products are based on the following considerations:

- 801 • There is at present no epidemiological evidence of CJD and vCJD transmission by urine-derived  
802 medicinal products.
- 803 • TSE infectivity in urine has been reported in some animal models.
- 804 • Abnormal PrP has been detected by different methods in 40% of sCJD patient urine samples  
805 and 93% of vCJD samples.

806       • The review of manufacturing processes is described below.

807       Urine should be collected from countries where there is a surveillance system for both human and  
808       animal TSEs unless otherwise justified. It is noted that urine-derived medicinal products are not  
809       sourced from urine collected in the UK. Based on the limited data on human exposure to BSE-risk  
810       materials in other countries, it is still difficult to estimate the epidemiological risk in those countries  
811       which have a small number of vCJD cases or may have a TSE exposure risk.

812       For particular products, such as hormones from a relatively small well-defined donor population, some  
813       manufacturers have put in place limited exclusion criteria for the selection of a donor for inclusion in a  
814       donor panel. For other products manufactured from very large donor pools (e.g. urokinase), such  
815       measures are more difficult to apply. The use of exclusion criteria for selection for a donor panel is  
816       encouraged. The same exclusion criteria should be applied with respect to CJD and vCJD as used for  
817       blood/plasma donors providing starting material for the manufacture of plasma-derived medicinal  
818       products. Manufacturers should follow up the donor criteria at defined intervals. The exclusion of  
819       donors with known inflammation of kidney and/or chronic renal inflammatory diseases is encouraged.

820       Manufacturers are required to estimate the potential of their specific manufacturing processes to  
821       reduce infectivity following the same general, stepwise approach as recommended for plasma derived  
822       medicinal products (see Section 9.2.3). Extrapolation of results for plasma-derived medicinal products  
823       is not justified particularly for chromatographic steps at the beginning of the manufacturing process  
824       because of the high protein content in plasma. Investigational studies of infectivity reduction by the  
825       manufacturing processes should address potential accumulation of infectivity/PrP<sup>TSE</sup> on  
826       chromatographic columns or a potential batch to batch contamination due to carry-over of  
827       infectivity/PrP<sup>TSE</sup>. For inactivation studies, investigation of different TSE strains should be considered as  
828       they may vary in resistance.

829       General review of the manufacturing processes indicates that, in each manufacturing process, there is  
830       at least one step that might be theoretically capable of reducing infectivity if it were present in the  
831       starting material. In cases where the reduction capacity is limited, manufacturers should consider the  
832       addition of steps that may increase the overall removal capacity.

833       Record keeping for traceability is recommended for products where it is possible to trace back to donor  
834       level.

835

## 836 References

- 837 1. Will RG, Ironside JW, Zeidler M, Cousens SN, Estbeiro K, Alperovitch A, Poser S, Pocchiari M,  
838 Hofman A, Smith PG. A new variant of Creutzfeldt-Jakob disease in the UK. *Lancet* 1996;  
839 47:921-25.
- 840 2. Bruce ME, Will RG, Ironside JW, McConnell I, Drummond D, Suttle A, McCardle L, Chree A, Hope  
841 J, Birkett C, Cousens S, Fraser H, Bostock CJ. Transmissions to mice indicate that "new variant"  
842 CJD is caused by the BSE agent. *Nature* 1997;389:498-501.
- 843 3. Hill AF, Desbruslais M, Joiner S, Sidle KCL, Gowland I, Collinge J. The same prion strain causes  
844 vCJD and BSE. *Nature* 1997;389:448-450.
- 845 4. Scott MR, Will R, Ironside J, Nguyen HO, Tremblay P, DeArmond SJ, Prusiner SB. Compelling  
846 transgenic evidence for transmission of bovine spongiform encephalopathy prions to humans.  
847 *Proc. Natl. Acad. Sci* 1999;96:15137-15142.
- 848 5. EMA
- 849 5a Guideline on the investigation of manufacturing processes for plasma-derived medicinal  
850 products with regard to vCJD risk. October 2004,  
851 EMEA/CPMP/BWP/5136/03.<http://www.ema.europa.eu/pdfs/human/bwp/513603en.pdf>
- 852 5b CPMP Position Statement on Creutzfeldt-Jakob disease and plasma-derived and urine-  
853 derived medicinal products. February 2003,  
854 EMEA/CPMP/BWP/2879/02.<http://www.ema.europa.eu/pdfs/human/press/pos/287902en.pdf>
- 855 5c Summary Report of EMEA Expert Meeting on Human TSEs and medicinal products derived  
856 from human blood and plasma, 1 December 2000, EMEA/CPMP/BWP/450/01, 28 March  
857 2001.<http://www.ema.europa.eu/pdfs/human/regaffair/045001en.pdf>
- 858 5d Report of EMEA Expert Workshop on Human TSEs and plasma-derived medicinal products,  
859 15-16 May 2000, EMEA/CPMP/BWP/1244/00, 26 July 2000.  
860 <http://www.ema.europa.eu/pdfs/human/regaffair/045001en.pdf>
- 861 5e EMEA Workshop on application to pharmaceuticals of assays for markers of TSE, February  
862 1999, CPMP/257/99.<http://www.ema.europa.eu/pdfs/human/bwp/025799en.pdf>
- 863 5f CPMP Position Statement on New variant CJD and plasma-derived medicinal products,  
864 February 1998, CPMP/201/98.<http://www.ema.europa.eu/pdfs/human/press/pos/020198en.pdf>
- 865 5g Report on Expert Workshop on CJD Risk and Urine-derived medicinal products EMEA,  
866 London - 12-13 July 2007. <http://www.ema.europa.eu/pdfs/human/bwp/47271709en.pdf>
- 867 5h CHMP position statement on Creutzfeldt-Jakob disease and plasma-derived and urine-  
868 derived medicinal products. 23 June 2011 EMA/CHMP/BWP/303353/2010  
869 [http://www.ema.europa.eu/docs/en\\_GB/document\\_library/Position\\_statement/2011/06/WC5001](http://www.ema.europa.eu/docs/en_GB/document_library/Position_statement/2011/06/WC5001)  
870 [08071.pdf](http://www.ema.europa.eu/docs/en_GB/document_library/Position_statement/2011/06/WC5001)
- 871 6. Llewelyn CA, Hewitt PE, Knight RSG, Amar K, Cousens S, Mackenzie J, Will RG. Possible  
872 transmission of variant Creutzfeldt-Jakob disease by blood transfusion. *Lancet* 2004;363:417-  
873 421.
- 874 7. European Commission



- 875 7a Directive 2002/98/EC of the European Parliament and Council setting standards of quality  
876 and safety for the collection, testing, processing, storage and distribution of human blood and  
877 blood components and amending Directive 2001/83 EC, OJ L 33, 8.2.2003, pp. 30-40.  
878 <http://eur-lex.europa.eu/LexUriServ/LexUriServ.do?uri=OJ:L:2003:033:0030:0040:EN:PDF>
- 879 7b Commission Directive 2004/33/EC implementing Directive 2002/98/EC of the European  
880 Parliament and of the Council as regards certain technical requirements for blood and blood  
881 components, OJ L 91,30.3.2004, pp.25-39. [http://eur-](http://eur-lex.europa.eu/LexUriServ/LexUriServ.do?uri=OJ:L:2004:091:0025:0039:EN:PDF)  
882 [lex.europa.eu/LexUriServ/LexUriServ.do?uri=OJ:L:2004:091:0025:0039:EN:PDF](http://eur-lex.europa.eu/LexUriServ/LexUriServ.do?uri=OJ:L:2004:091:0025:0039:EN:PDF)
- 883 7c Council Recommendation of 29 June 1998 on the suitability of blood and plasma donors  
884 and the screening of donated blood in the European Community (98/463/EC), OJ L 203 pp.14-26.  
885 <http://eur-lex.europa.eu/LexUriServ/LexUriServ.do?uri=OJ:L:1998:203:0014:0026:EN:PDF>
- 886 7d Statement of Commissioner Byrne on possible vCJD transmission by blood. Brussels, 18  
887 December 2003, IP/03/1781.  
888 [http://europa.eu/rapid/pressReleasesAction.do?reference=IP/03/1781&format=HTML&aged=0&i](http://europa.eu/rapid/pressReleasesAction.do?reference=IP/03/1781&format=HTML&aged=0&language=EN&guiLanguage=en)  
889 [anguage=EN&guiLanguage=en](http://europa.eu/rapid/pressReleasesAction.do?reference=IP/03/1781&format=HTML&aged=0&language=EN&guiLanguage=en)
- 890 7e Summary Statement of Technical Meeting of Blood Experts related to vCJD transmission.  
891 Luxembourg 20 January 2004, D(2004) FMD/360007.
- 892 8. "Guide to the preparation, use and quality assurance of blood components", Recommendation  
893 No. R (95) 16, 16th ed. 2011. Council of Europe.
- 894 9. European Commission: SCMPMD, SSC, SCENIHR  
895 SCMPMD
- 896 9a Opinion on the Safety of Human-Derived Products with Regard to TSE's, adopted on 18  
897 January 2002. [http://ec.europa.eu/health/ph\\_risk/committees/scmp/documents/out40\\_en.pdf](http://ec.europa.eu/health/ph_risk/committees/scmp/documents/out40_en.pdf)
- 898 9b Opinion on update of the opinion on the Risk Quantification for CJD Transmission via  
899 Substances of Human Origin, adopted on 16 February 2000.  
900 [http://ec.europa.eu/health/ph\\_risk/committees/scmp/documents/out28\\_en.pdf](http://ec.europa.eu/health/ph_risk/committees/scmp/documents/out28_en.pdf)
- 901 9c Opinion on the Policy Regarding the Use of Blood and Blood Products adopted by Written  
902 Procedure on 24 March 1999.  
903 [http://ec.europa.eu/health/scientific\\_committees/emerging/opinions/scmpmd/scmp\\_out20\\_en.h](http://ec.europa.eu/health/scientific_committees/emerging/opinions/scmpmd/scmp_out20_en.htm)  
904 [tm](http://ec.europa.eu/health/scientific_committees/emerging/opinions/scmpmd/scmp_out20_en.htm)
- 905 9d Opinion on the risk quantification for CJD transmission via substances of human origin,  
906 adopted on 21/10/98.  
907 [http://ec.europa.eu/health/ph\\_risk/committees/scmp/documents/out12\\_en.pdf](http://ec.europa.eu/health/ph_risk/committees/scmp/documents/out12_en.pdf)
- 908 SSC
- 909 9e Opinion on the Implications of the Recent Papers on Transmission of BSE by Blood  
910 Transfusion in Sheep (Houston et al, 2000; Hunter et al, 2002), adopted September 2002.  
911 [http://ec.europa.eu/food/fs/sc/ssc/out79\\_en.pdf](http://ec.europa.eu/food/fs/sc/ssc/out79_en.pdf)
- 912 9f Opinion on the Implications of the Houston et al paper in The Lancet of 16 September  
913 2000 on the Transmission of BSE by blood transfusion in sheep. (The Lancet, Vol.356, pp999-  
914 1000;955-956;1013). [http://ec.europa.eu/food/fs/sc/ssc/out143\\_en.pdf](http://ec.europa.eu/food/fs/sc/ssc/out143_en.pdf)

- 915 9g Oral exposure of Humans to the BSE agent: infective dose and species barrier adopted by  
916 the SSC at its meeting of 13-14 April 2000 following a public consultation via Internet between 6  
917 and 27 March 2000. [http://ec.europa.eu/food/fs/sc/ssc/out79\\_en.pdf](http://ec.europa.eu/food/fs/sc/ssc/out79_en.pdf)
- 918 9h Opinion on the Human Exposure Risk (HER) via food with respect to BSE - Adopted on 10  
919 December 1999. [http://ec.europa.eu/food/fs/sc/ssc/out67\\_en.pdf](http://ec.europa.eu/food/fs/sc/ssc/out67_en.pdf)
- 920 SCENIHR
- 921 9i Opinion on the Safety of Human-derived Products with regard to Variant Creutzfeldt-Jakob  
922 Disease adopted on 11-12 May 2006.  
923 [http://ec.europa.eu/health/archive/ph\\_risk/committees/04\\_scenihr/docs/scenihr\\_o\\_004b.pdf](http://ec.europa.eu/health/archive/ph_risk/committees/04_scenihr/docs/scenihr_o_004b.pdf)
- 924 10. WHO
- 925 10a WHO Guidelines on Transmissible Spongiform Encephalopathies in relation to Biological  
926 and Pharmaceutical Products. 2003. <http://www.who.int/biologicals>
- 927 10b WHO Guidelines on Tissue Infectivity Distribution in Transmissible Spongiform  
928 Encephalopathies. 2006. <http://www.who.int/biologicals>
- 929 10c WHO Tables on Tissue Infectivity Distribution in Transmissible Spongiform  
930 Encephalopathies. Updated 2010. <http://www.who.int/bloodproducts/tablestissueinfectivity.pdf>
- 931 11. Council of Europe Recommendation Rec(2001)4 on the prevention of the possible transmission  
932 of variant Creutzfeldt-Jakob Disease (vCJD) by blood  
933 transfusion. [http://www.coe.int/t/dg3/health/recommendations\\_en.asp](http://www.coe.int/t/dg3/health/recommendations_en.asp)
- 934 12. Ladogana A, Puopolo M, Croes EA, Budka H, Jarius C, Collins S, Klug GM, Sutcliffe T, Giulivi A,  
935 Alperovitch A, Delasnerie-Laupretre N, Brandel JP, Poser S, Kretzschmar H, Rietveld I, Mitrova E,  
936 Cuesta Jde P, Martinez-Martin P, Glatzel M, Aguzzi A, Knight R, Ward H, Pocchiari M, van Duyn  
937 CM, Will RG, Zerr I. Mortality from Creutzfeldt-Jakob disease and related disorders in Europe,  
938 Australia, and Canada. *Neurology*. 2005;64:1586-91.
- 939 13. Dorsey K, Zou S, Schonberger LB, Sullivan M, Kessler D, Notari E 4th, Fang CT, Dodd RY. Lack of  
940 evidence of transfusion transmission of Creutzfeldt-Jakob disease in a US surveillance study.  
941 *Transfusion* 2009;49: 977-984.
- 942 14. Urwin P, Thanigaikumar K, Ironside JW, Molesworth A, Knight RS, Hewitt PE, Llewelyn C,  
943 Mackenzie J, Will RG. Sporadic Creutzfeldt-Jakob Disease in 2 Plasma Product Recipients, United  
944 Kingdom. *Em Inf Dis*. 2017;6: 893-7.
- 945 15. Diack AB, Ritchie DL, Peden AH, Brown D, Boyle A, Morabito L, Maclennan D, Burgoyne P, Jansen  
946 C, Knight RS, Piccardo P, Ironside JW, Manson JC. Variably protease-sensitive prionopathy, a  
947 unique prion variant with inefficient transmission properties. *Emerg Infect Dis*. 2014;20:1969-  
948 79.
- 949 16. Notari S, Xiao X, Espinosa JC, Cohen Y, Qing L, Aguilar-Calvo P, Kofskey D, Cali I, Cracco L,  
950 Kong Q, Torres JM, Zou W, Gambetti P. Transmission characteristics of variably protease-  
951 sensitive prionopathy. *Emerg Infect Dis*. 2014;20:2006-14.
- 952 17. UK Monthly Creutzfeldt Jakob disease statistics.  
953 <https://www.cjd.ed.ac.uk/>

- 954 18. French Monthly Creutzfeldt Jakob disease statistics.  
955 [http://invs.santepubliquefrance.fr/Dossiers-thematiques/Maladies-infectieuses/Risques-  
infectieux-d-origine-alimentaire/Maladie-de-Creutzfeldt-Jakob/Donnees-epidemiologiques](http://invs.santepubliquefrance.fr/Dossiers-thematiques/Maladies-infectieuses/Risques-<br/>956 infectieux-d-origine-alimentaire/Maladie-de-Creutzfeldt-Jakob/Donnees-epidemiologiques)
- 957 19. Worldwide vCJD statistics.  
958 <https://www.cjd.ed.ac.uk/data-and-reports/variant-cjd-cases-worldwide>.
- 959 20. Ward HJ, MacKenzie JM, Llewelyn CA, Knight RS, Hewitt PE, Connor N, Molesworth A, Will RG.  
960 Variant Creutzfeldt-Jakob disease and exposure to fractionated plasma products. *Vox Sang*.  
961 2009;97:207-10.
- 962 21. Mok T, Jaunmuktane Z, Joiner S, Campbell T, Morgan C, Wakerley B, Golestani F, Rudge P, Mead  
963 S, Jäger HR, Wadsworth JD, Brandner S, Collinge J. Variant Creutzfeldt-Jakob Disease in a  
964 Patient with Heterozygosity at PRNP Codon 129. *NEJM* 2017;376:292-294.
- 965 22. Istituto Superiore di Sanità  
966 <http://www.epicentro.iss.it/problemi/jakob/EpidemiologiciItalia.asp>
- 967 23. Hilton DA, Ghani AC, Conyers L, Edwards P, McCardle L, Ritchie D, Penney M, Hegazy D, Ironside  
968 JW. Prevalence of lymphoreticular prion protein accumulation in UK tissue samples. *Journal of*  
969 *Pathology* 2004;203:733-739.
- 970 24. Ironside JW, Bishop MT, Connolly K, Hegazy D, Lowrie S, Le Grice M, Ritchie DL, McCardle LM,  
971 Hilton DA. Variant Creutzfeldt-Jakob disease: prion protein genotype analysis of positive  
972 appendix tissue samples from a retrospective prevalence study. *BMJ*. 2006;332:1186-8.
- 973 25. Ghani AC, Donnelly CA, Ferguson NM, Anderson RM. Updated projections of future vCJD deaths  
974 in the UK. *BMC Infect Dis* 2003;3:4.
- 975 26. Clewley JP, Lelly CM, Andrews N, Vogliqi K, Mallinson G, Kaiser M, Hilton DA, Ironside JW,  
976 Edwards P, McCardle LM, Ritchie DL, Dabaghian, R, Ambrose HE, Gill ON. Prevalence of disease-  
977 related prion protein in anonymous tonsil specimens in Britain: cross sectional opportunistic  
978 study. *BMJ* 2009;338:b1442.
- 979 27. De Marco MF, Linehan J, Gill ON, Clewley JP, Brandner S. Large-scale immunohistochemical  
980 examination for lymphoreticular prion protein in tonsil specimens collected in Britain. *J Pathol*  
981 2010;222:380-387.
- 982 28. Gill ON, Spencer Y, Richard-Loendt A, Kelly C, Dabaghian R, Boyes L, Linehan J, Simmons M,  
983 Webb P, Bellerby P, Andrews N, Hilton DA, Ironside JW, Beck J, Poulter M, Mead S, Brandner  
984 S. Prevalent abnormal prion protein in human appendixes after bovine spongiform  
985 encephalopathy epizootic: large scale survey. *BMJ*. 2013 Oct;347:f5675.
- 986 29. Summary results of the third national survey of abnormal prion prevalence in archived appendix  
987 specimens. In *Health Protection Report* Vol. 10 No. 26 – 12 August  
988 2016. <https://www.gov.uk/government/publications/health-protection-report-volume-10-2016>.
- 989 30. Balkema-Buschmann A, Eiden M, Hoffmann C, Kaatz M, Ziegler U, Keller M Groschup MH. BSE  
990 infectivity in the absence of detectable PrPSc accumulation in the tongue and nasal mucosa of  
991 terminally diseased cattle. *J Gen Virol* 2011;92:467–476.
- 992 31. Piccardo P, Manson J, King D, Ghetti B, Barron RM. Accumulation of prion protein in the brain  
993 that is not associated with transmissible disease. *PNAS* 2007;104:4712-4717.



- 994 32. Tixador P, Herzog L, Reine F, Jaumain E, Chapuis J, Le Dur A, Laude H, Beringue V. The physical  
995 relationship between infectivity and prion protein aggregates is strain-dependent. *PLoS Pathog.*  
996 2010;6:e1000859.
- 997 33. Safar JG, Geschwind MD, Deering C, Didorenko S, Sattavat M, Sanchez H, Serban A, Vey M,  
998 Baron H, Giles K, Miller BL, Dearmond SJ, Prusiner SB. Diagnosis of human prion disease. *PNAS*  
999 2005; 102:3501–3506.
- 1000 34. Choi YP, Groner A, Ironside JW, Head MW. Comparison of the level, distribution and form of  
1001 disease-associated prion protein in variant and sporadic Creutzfeldt–Jakob diseased brain using  
1002 conformation-dependent immunoassay and Western blot. *J Gen Virol* 2011;92:727–732.
- 1003 35. Wadsworth JD, Joiner S, Hill AF, Campbell TA, Desbruslais M, Luthert PJ, Collinge J. Tissue  
1004 distribution of protease resistant prion protein in variant Creutzfeldt-Jakob disease using a highly  
1005 sensitive immunoblotting assay. *Lancet* 2001;358:171-180.
- 1006 36. Head MW, Ritchie D, Smith N, McLoughlin V, Nailon W, Samad S, Masson S, Bishop M, McCardle  
1007 L, Ironside JW. Peripheral tissue involvement in sporadic, iatrogenic, and variant Creutzfeldt-  
1008 Jakob disease: an immunohistochemical, quantitative, and biochemical study. *Am J Pathol.* 2004  
1009 Jan;164(1):143-53.. *American Journal of Pathology* 2004;164:143-153.
- 1010 37. Glatzel M, Abela E, Maissen M, Aguzzi A. Extraneural pathologic prion protein in sporadic  
1011 Creutzfeldt-Jakob disease. *New England Journal of Medicine* 2003;349:1812-1820.
- 1012 38. Mead S, Wadsworth JF, Porter M, Linehan JM, Pietkiewicz W, Jackson GS, Brandner S, Collinge J.  
1013 Variant Creutzfeldt-Jakob disease with extremely low lymphoreticular deposition of prion protein.  
1014 *JAMA Neurol* 2014;71:340–343.
- 1015 39. Rubenstein R, Chang B. Re-Assessment of PrPSc Distribution in Sporadic and Variant CJD. *PLoS*  
1016 *ONE* 2013;8:e66352.
- 1017 40. Bruce ME, McConnell I, Will RG, Ironside JW. Detection of variant Creutzfeldt-Jakob disease  
1018 infectivity in extraneural tissues. *Lancet* 2001;358:208-209.
- 1019 41. Accelerated, spleen-based titration of variant Creutzfeldt-Jakob disease infectivity in transgenic  
1020 mice expressing human prion protein with sensitivity comparable to that of survival time  
1021 bioassay. Halliez S, Reine F, Herzog L, Jaumain E, Haik S, Rezaei H, Vilotte JL, Laude H,  
1022 Beringue V. *J Virol.* 2014;88:8678-86.
- 1023 42. Bishop MT, Diack AB, Ritchie DL, Ironside JW, Will RG, Manson JC. Prion infectivity in the spleen  
1024 of a *PRNP* heterozygous individual with subclinical variant Creutzfeldt–Jakob disease. *Brain*  
1025 2013;136:1139–1145.
- 1026 43. Peden AH, Ritchie DL, Head MW, Ironside JW. Detection and localization of PrPSc in the skeletal  
1027 muscle of patients with variant, iatrogenic, and sporadic forms of Creutzfeldt-Jakob disease. *Am*  
1028 *J Pathol.* 2006;168:927-935.
- 1029 44. Brown P, Rohwer RG, Dunstan BC, MacAuley C, Gajdusek DC, Drohan WN. The distribution of  
1030 infectivity in blood components and plasma derivatives in experimental models of transmissible  
1031 spongiform encephalopathy. *Transfusion* 1998;38:810-816.
- 1032 45. Cervenakova L, Yakovleva O, McKenzie C, Kolchinsky S, McShane L, Drohan WN, Brown P.  
1033 Similar levels of infectivity in the blood of mice infected with human-derived vCJD and GSS  
1034 strains of transmissible spongiform encephalopathy. *Transfusion* 2003;43:1687-1694.

- 1035 46. Brown P, Cervenáková L, McShane LM, Barber P, Rubenstein R, Drohan WN. Further studies of  
1036 blood infectivity in an experimental model of transmissible spongiform encephalopathy, with an  
1037 explanation of why blood components do not transmit Creutzfeldt-Jakob disease in humans.  
1038 *Transfusion* 1999;39:1169-1178.
- 1039 47. Gregori L, Lambert BC, Gurgel PV, Gheorghiu L, Edwardson P, Lathrop JT, MacAuley C, Carbonell  
1040 RG, Burton SJ, Hammond D, Rohwer RG. Reduction of transmissible spongiform encephalopathy  
1041 infectivity from human red blood cells with prion affinity ligands. *Tranfusion* 2006; 46: 1152-  
1042 1161.
- 1043 48. Lefrère JJ, Hewitt P. From mad cows to sensible blood transfusion: the risk of prion transmission  
1044 by labile blood components in the United Kingdom and in France. *Transfusion* 2009;49:797-812.
- 1045 49. Holada K, Vostal JG, Theisen PW, MacAuley C, Gregori L, Rohwer RG. Scrapie infectivity in  
1046 hamster blood is not associated with platelets. *J Virol.* 2002 76:4649-4650.
- 1047 50. Bons N, Lehmann S, Mestre-Francès N, Dormont D, Brown P. Brain and buffy coat transmission  
1048 of bovine spongiform encephalopathy to the primate *Microcebus murinus*. *Transfusion*  
1049 2002;42:513-516.
- 1050 51. Lescoutra-Etcheagaray N, Jaffré N, Sumian C, Durand V, Correia E, Mikol J, Luccantoni-Freire S,  
1051 Culeux A, Deslys JP, Comoy EE. Evaluation of the protection of primates transfused with variant  
1052 Creutzfeldt-Jakob disease-infected blood products filtered with prion removal devices: a 5-year  
1053 update. *Transfusion.* 2015;1231-41.
- 1054 52. Lacroux C, Vilette D, Fernández-Borges N, Litaise C, Lugan S, Morel N, Corbière F, Simon S,  
1055 Simmons H, Costes P, Weisbecker JL, Lantier I, Lantier F, Schelcher F, Grassi J, Castilla J,  
1056 Andréoletti O. Prionemia and leukocyte-platelet-associated infectivity in sheep transmissible  
1057 spongiform encephalopathy models. *J Virol.* 2012;86:2056-66.
- 1058 53. Andréoletti O, Litaise C, Simmons H, Corbière F, Lugan S, Costes P, Schelcher F, Vilette D,  
1059 Grassi J, Lacroux C. Highly efficient prion transmission by blood transfusion. *PloS Pathog.* 2012;  
1060 8(6):e1002782.
- 1061 54. Mathiason CK, Hayes-Klug J, Hays SA, Powers J, Osborn DA, Dahmes SJ, Miller KV, Warren RJ,  
1062 Mason GL, Telling GC, Young AJ, Hoover EA. B cells and platelets harbor prion infectivity in the  
1063 blood of deer infected with chronic wasting disease. *J Virol.* 2010;84:5097-5107.
- 1064 55. Ironside JW. Variant Creutzfeldt-Jakob disease: risk of transmission by blood transfusion and  
1065 blood therapies. *Haemophilia.* 2006; Suppl 1:8-15.
- 1066 56. Houston F, Foster JD, Chong A, Hunter N, Bostock CJ. Transmission of BSE by blood transfusion  
1067 in sheep. *Lancet* 2000;356:999-1000.
- 1068 57. Hunter N, Foster J, Chong A, McCutcheon S, Parnham D, Eaton S, MacKenzie C, Houston F.  
1069 Transmission of prion diseases by blood transfusion. *Journal of General Virology* 2002;83:2897-  
1070 2905.
- 1071 58. Lacroux C, Bougard D, Litaise C, Simmons H, Corbiere F, Dernis D, Tardivel R, Morel N, Simon S,  
1072 Lugan S, Costes P, Weisbecker JL, Schelcher F, Grassi J, Coste J, Andréoletti O. Impact of  
1073 leucocyte depletion and prion reduction filters on TSE blood borne transmission. *PLoS One.*  
1074 2012;7(7):e42019.

- 1075 59. McCutcheon S, Alejo Blanco AR, Houston EF, de Wolf C, Tan BC, Smith A, Groschup MH, Hunter  
1076 N, Hornsey VS, MacGregor IR, Prowse CV, Turner M, Manson JC. All clinically-relevant blood  
1077 components transmit prion disease following a single blood transfusion: a sheep model of vCJD.  
1078 PLoS One. 2011;6(8):e23169.
- 1079 60. Gregori L, Yang H, Anderson S. Estimation of variant Creutzfeldt-Jakob disease infectivity titers  
1080 in human blood. *Transfusion*. 2011;51:2596-602.
- 1081 61. Douet JY, Lacroux C, Litaize C, Lugan S, Corbière F, Arnold M, Simmons H, Aron N, Costes P,  
1082 Tillier C, Cassard H, Andréoletti O. Mononucleated Blood Cell Populations Display Different  
1083 Abilities To Transmit Prion Disease by the Transfusion Route. *J Virol*. 2016;90:3439-45.
- 1084 62. Castilla J, Saá P, Soto C. Detection of prions in blood. *Nat Med*. 2005;11:982-5.
- 1085 63. Saá P, Yakovleva O, de Castro J, Vasilyeva I, De Paoli SH, Simak J, Cervenakova L. First  
1086 demonstration of transmissible spongiform encephalopathy-associated prion protein (PrP<sup>Sc</sup>) in  
1087 extracellular vesicles from plasma of mice infected with mouse-adapted variant Creutzfeldt-  
1088 Jakob disease by in vitro amplification. *J Biol Chem*. 2014;289:29247-60.
- 1089 64. Elder AM, Henderson DM, Nalls AV, Hoover EA, Kincaid AE, Bartz JC, Mathiason CK. Immediate  
1090 and Ongoing Detection of Prions in the Blood of Hamsters and Deer following Oral, Nasal, or  
1091 Blood Inoculations. *J Virol*. 2015;89:7421-4.
- 1092 65. Sawyer EB, Edgeworth JA, Thomas C, Collinge J, Jackson GS. Preclinical detection of infectivity  
1093 and disease-specific PrP in blood throughout the incubation period of prion disease. *Sci Rep*.  
1094 2015; 5:17742.
- 1095 66. Properzi F, Logozzi M, Abdel-Haq H, Federici C, Lugini L, Azzarito T, Cristofaro I, di Sevo D,  
1096 Ferroni E, Cardone F, Venditti M, Colone M, Comoy E, Durand V, Fais S, Pocchiari M. Detection of  
1097 exosomal prions in blood by immunochemistry techniques. *J Gen Virol*. 2015;96:1969-74.
- 1098 67. Lacroux C, Comoy E, Moudjou M, Perret-Liaudet A, Lugan S, Litaize C, Simmons H, Jas-Duval C,  
1099 Lantier I, Béringue V, Groschup M, Fichet G, Costes P, Streichenberger N, Lantier F, Deslys JP,  
1100 Vilette D, Andréoletti O. Preclinical detection of variant CJD and BSE prions in blood. *PLoS*  
1101 *Pathog*. 2014;10:e1004202.
- 1102 68. Halliez S, Jaumain E, Huor A, Douet JY, Lugan S, Cassard H, Lacroux C, Béringue V, Andréoletti  
1103 O, Vilette D. White blood cell-based detection of asymptomatic scrapie infection by ex vivo  
1104 assays. *PLoS One*. 2014;9:e104287.
- 1105 69. Bannach O, Birkmann E, Reinartz E, Jaeger KE, Langeveld JP, Rohwer RG, Gregori L, Terry LA,  
1106 Willbold D, Riesner D. Detection of prion protein particles in blood plasma of scrapie infected  
1107 sheep. *PLoS One*. 2012;7:e36620.
- 1108 70. McDowell KL, Nag N, Franco Z, Bu M, Piccardo P, Cervenak J, Deslys JP, Comoy E, Asher DM,  
1109 Gregori L. Blood reference materials from macaques infected with variant Creutzfeldt-Jakob  
1110 disease agent. *Transfusion*. 2015;55:405-12.
- 1111 71. Murayama Y, Masujin K, Imamura M, Ono F, Shibata H, Tobiume M, Yamamura T, Shimozaki N,  
1112 Terao K, Yamakawa Y, Sata T. Ultrasensitive detection of PrP(Sc) in the cerebrospinal fluid and  
1113 blood of macaques infected with bovine spongiform encephalopathy prion. *J Gen Virol*.  
1114 2014;95:2576-88.



- 1115 72. Gillies M, Chohan G, Llewelyn CA, MacKenzie J, Ward HJ, Hewitt PE, Will RG. A retrospective case  
1116 note review of deceased recipients of vCJD-implicated blood transfusions. *Vox Sang* 97: 211-  
1117 218, 2009.
- 1118 73. Peden AH, Head MW, Ritchie DL, Bell JE, Ironside JW. Preclinical vCJD after blood transfusion in  
1119 a PRNP codon 129 heterozygous patient. *Lancet*. 2004, 7-13;364:527-9
- 1120 74. Bishop MT, Diack AB, Ritchie DL, Ironside JW, Will RG, Manson JC. Prion infectivity in the spleen  
1121 of a PRNP heterozygous individual with subclinical variant Creutzfeldt-Jakob disease. *Brain*.  
1122 2013;136:1139-45.
- 1123 75. Peden A, McCardle L, Head MW, Love S, Ward HJ, Cousens SN, Keeling DM, Millar CM, Hill FG,  
1124 Ironside JW. Variant CJD infection in the spleen of a neurologically asymptomatic UK adult  
1125 patient with haemophilia. *Haemophilia* 2010, 16:296-304.
- 1126 76. Douet JY, Zafar S, Perret-Liaudet A, Lacroux C, Lugan S, Aron N, Cassard H, Ponto C, Corbière F,  
1127 Torres JM, Zerr I, Andreoletti O. Detection of infectivity in blood of persons with variant and  
1128 sporadic Creutzfeldt-Jakob disease. *Emerg Infect Dis*. 2014;20:114-7.
- 1129 77. Urwin PJM, Mackenzie JM, Llewelyn CA, Will RG, Hewitt PE. Creutzfeldt–Jakob disease and blood  
1130 transfusion: updated results of the UK Transfusion Medicine Epidemiology Review Study. *Vox*  
1131 *Sanguinis*. 2016;110:310–316.
- 1132 78. Crowder LA, Schonberger LB, Dodd RY, Steele WR. Creutzfeldt-Jakob disease lookback study: 21  
1133 years of surveillance for transfusion transmission risk. *Transfusion*. 2017;57:1875-1878.
- 1134 79. Manuelidis EE, Kim JH, Mericangas JR, Manuelidis L. Transmission to animals of Creutzfeldt-  
1135 Jakob disease from human blood. *Lancet* 2: 896-7, 1985.
- 1136 80. Tateishi J. Transmission of Creutzfeldt-Jakob disease from human blood and urine into mice.  
1137 *Lancet* 2: 1074, 1985.
- 1138 81. Tamai Y, Kojima H, Kitajima R, Taguchi F, Ohtani Y, Kawaguchi T, Miura S, Sato M, Ishihara Y.  
1139 Demonstration of the transmissible agent in tissue from a pregnant woman with Creutzfeldt-  
1140 Jakob disease. *N Engl J Med* 327: 649, 1992.
- 1141 82. Deslys JP, Lasmézas C, Dormont D. Selection of specific strains in iatrogenic Creutzfeldt-Jakob  
1142 disease. *Lancet* 343: 848-849, 1994.
- 1143 83. Cervenakova L, Brown P, Hammond DJ, Lee CA, Saenko EL. Review article: Factor VIII and  
1144 transmissible spongiform encephalopathy: the case for safety. *Haemophilia* 2002;8:63-75.
- 1145 84. Brown P, Gibbs CJ Jr, Rodgers-Johnson P, Asher DM, Sulima MP, Bacote A, Goldfarb LG,  
1146 Gajdusek DC. Human spongiform encephalopathy: the National Institutes of Health series of 300  
1147 cases of experimentally transmitted disease. *Ann Neurol* 35: 513-529, 1994.
- 1148 85. Edgeworth JA, Farmer M, Sicilia A, Tavares P, Beck J, Campbell T, Lowe J, Mead S, Rudge P,  
1149 Collinge J, Jackson GS. Detection of prion infection in variant Creutzfeldt-Jakob disease: a blood-  
1150 based assay. *Lancet* 2011, 377:487-493.
- 1151 86. Transmissible Spongiform Encephalopathy Advisory Committee June 12th 2009.  
1152 [www.fda.gov/TransmissibleSpongiformEncephalopathyAdvisory Committee](http://www.fda.gov/TransmissibleSpongiformEncephalopathyAdvisoryCommittee).
- 1153 87. Press Release. Amorfix Life Sciences Ltd. 29 December  
1154 2009. [http://www.amorfix.com/pdf\\_press/pr\\_2009/2009\\_12\\_29\\_Corporate\\_Update\\_Final.pdf](http://www.amorfix.com/pdf_press/pr_2009/2009_12_29_Corporate_Update_Final.pdf).

- 1155 88. Miele G, Manson J, Clinton M: A novel erythroid-specific marker of transmissible spongiform  
1156 encephalopathies. *Nat.Med.* 2001, 7:361-364.
- 1157 89. Feng L, Gell DA, Zhou S, Gu L, Kong Y, Li J, Hu M, Yan N, Lee C, Rich AM, Armstrong RS, Lay PA,  
1158 Gow AJ, Weiss MJ, Mackay JP, Shi Y. Molecular mechanism of AHSP-mediated stabilization of  
1159 alpha-hemoglobin. *Cell* 2004, 119:629-640.
- 1160 90. Cepek L, Brechlin P, Steinacker P, Mollenhauer B, Klingebiel E, Bibl M, Kretzschmar HA, Wiltfang  
1161 J, Otto M. Proteomic Analysis of the Cerebrospinal Fluid of Patients with Creutzfeldt-Jakob  
1162 Disease. *Dement.Geriatr.Cogn Disord.* 2006;23:22-28.
- 1163 91. Chiang YW, Otoshima Y, Watanabe Y, Inanami O, Shimoyama Y. Dynamics and Local Ordering of  
1164 Spin-Labeled Prion Protein: An ESR Simulation Study of a Highly PH-Sensitive Site. *J Biomol*  
1165 *Struct Dyn* 2008, 26:355-366.
- 1166 92. Brown AR, Blanco AR, Miele G, Hawkins SA, Hopkins J, Fazakerley JK, Manson J, Clinton M.  
1167 Differential expression of erythroid genes in prion disease. *Biochem Biophys Res Commun* 2007,  
1168 364:366-371.
- 1169 93. Quattieri A, Urso E, Pera ML, Sprovieri T, Bossio S, Gambardella A, Quattrone A. Proteomic  
1170 profiling of cerebrospinal fluid in Creutzfeldt-Jakob disease. *Expert Rev Proteomics* 2010, 7:907-  
1171 917.
- 1172 94. Appleford NE, Wilson K, Houston F, Bruce LJ, Morrison A, Bishop M, Chalmers K, Miele G, Massey  
1173 E, Prowse C, Manson J, Will RG, Clinton M, MacGregor I, Anstee DJ. Alpha-Hemoglobin stabilizing  
1174 protein is not a suitable marker for a screening test for variant Creutzfeldt-Jakob disease.  
1175 *Transfusion* 2008,48:1616-1626.
- 1176 95. Atarashi R, Moore RA, Sim VL, Hughson AG, Dorward DW, Onwubiko HA, Priola SA, Caughey B:  
1177 Ultrasensitive detection of scrapie prion protein using seeded conversion of recombinant prion  
1178 protein. *Nat Methods* 2007, 4:645-650.
- 1179 96. Saborio GP, Permanne B, Soto C: Sensitive detection of pathological prion protein by cyclic  
1180 amplification of protein misfolding. *Nature* 2001, 411:810-813.
- 1181 97. McGuire LI, Poggioli A, Poggiolini I, Suardi S, Grznarova K, Shi S, De Vil B, Sarros S, Satoh K,  
1182 Cheng K, Cramm M, Fairfoul G, Schmitz M, Zerr I, Cras P, Equestre M, Tagliavini F, Atarashi R,  
1183 Knox D, Collins S, Haik S, Parchi P, Pocchiari M, Green A: CSF RT-QuIC is a robust and reliable  
1184 test for sporadic CJD: An international study. *Ann Neurol* 2016;80:160-5.
- 1185 98. Moda F, Gambetti P, Notari S, Concha-Marambio L, Catania M, Park KW, Maderna E, Suardi S,  
1186 Haik S, Brandel JP, Ironside J, Knight R, Tagliavini F, Soto C: Prions in the urine of patients with  
1187 variant Creutzfeldt-Jakob disease. *N Engl J Med* 2014;371:530-539.
- 1188 99. Concha-Marambio, L., Pritzkow, S., Moda, F., Tagliavini, F., Ironside, J. W., Schulz, P. E., & Soto,  
1189 C. (2016). Detection of prions in blood from patients with variant Creutzfeldt-Jakob disease.  
1190 *Science Translational Medicine*, 8(370), [370ra183]. DOI: [10.1126/scitranslmed.aaf6188](https://doi.org/10.1126/scitranslmed.aaf6188)
- 1191 100. Bougard D et al. Detection of prions in the plasma of presymptomatic and symptomatic patients  
1192 with variant Creutzfeldt-Jakob disease. *Sci Transl Med* 2016 Dec 21;  
1193 8:370ra182. <http://dx.doi.org/10.1126/scitranslmed.aag1257>

- 1194 101. Guideline for the CE marking of blood based in vitro diagnostic medical devices for vCJD based  
 1195 on detection of abnormal PrP. European commission DG Health and Consumer Directorate.  
 1196 2012. [http://ec.europa.eu/health/medical-devices/files/meddev/2\\_14\\_4\\_ol\\_en.pdf](http://ec.europa.eu/health/medical-devices/files/meddev/2_14_4_ol_en.pdf).
- 1197 102. Jackson GS, Burk-Rafel J, Edgeworth JA, Sicilia A, Abdilahi S, Korteweg J, Mackey J, Thomas C,  
 1198 Wang G, Schott JM, Mummery C, Chinnery PF, Mead S, Collinge J. Population Screening for  
 1199 Variant Creutzfeldt-Jakob Disease Using a Novel Blood Test: Diagnostic Accuracy and Feasibility  
 1200 Study. *JAMA Neurol* 2014, 71:340.
- 1201 103. Wroe SJ, Pal S, Siddique D, Hyare H, Macfarlane R, Joiner S, Linehan J, Brandner S, Wadsworth  
 1202 JD, Hewitt P, Collinge J: Clinical presentation and pre-mortem diagnosis of variant Creutzfeldt-  
 1203 Jakob disease associated with blood transfusion: a case report. *Lancet* 2006, 368:2061-2067.
- 1204 104. Hewitt PE, Llewelyn CA, Mackenzie J, Will RG: Creutzfeldt-Jakob disease and blood transfusion:  
 1205 results of the UK Transfusion Medicine Epidemiological Review study. *Vox Sang.* 2006, 91:221-  
 1206 230.
- 1207 105. Cohen H, Bolton-Maggs P: Breaking records in blood transfusion safety. *Transfus.Med* 2012,  
 1208 22:241-243.
- 1209 106. Krailadsiri P, Seghatchian J, Macgregor I, Drummond O, Perrin R, Spring F, Prescott R,  
 1210 Williamson L, Prowse C, Anstee D, Turner M. The effects of leukodepletion on the generation and  
 1211 removal of microvesicles and prion protein in blood components. *Transfusion* 2006;46:407-17.
- 1212 107. Gregori L, Gurgel PV, Lathrop JT, Edwardson P, Lambert BC, Carbonell RG, Burton SJ, Hammond  
 1213 DJ, Rohwer RG. Reduction in infectivity of endogenous transmissible spongiform  
 1214 encephalopathies present in blood by adsorption to selective affinity resins. *Lancet* 2006; 368:  
 1215 2226-2230.
- 1216 108. SaBTO minutes meeting 27 Oct  
 1217 2009. [http://www.dh.gov.uk/prod\\_consum\\_dh/groups/dh\\_digitalassets/@dh/@ab/documents/di](http://www.dh.gov.uk/prod_consum_dh/groups/dh_digitalassets/@dh/@ab/documents/digitalasset/dh_108860.pdf)  
 1218 [gitalasset/dh\\_108860.pdf](http://www.dh.gov.uk/prod_consum_dh/groups/dh_digitalassets/@dh/@ab/documents/digitalasset/dh_108860.pdf)
- 1219 109. McLeod NP, Nugent P, Dixon D, Dennis M, Cornwall M, Mallinson G, Watkins N, Thomas S, Sutton  
 1220 JM. Evaluation of efficacy of prion reduction filters using blood from an endogenously infected  
 1221 263K scrapie hamster model. *Transfusion* 2015;55:2390-7.
- 1222 110. McCutcheon S, Alejo Blanco AR, Tan BC, Gonzalez L, Martin S, Mallinson G, Appleford NE, Turner  
 1223 ML, Manson JC, Houston EF: A prion reduction filter does not completely remove endogenous  
 1224 prion infectivity from sheep blood. *Transfusion* 2015;55:2123-33.
- 1225 111. Neisser-Svae A, Bailey A, Gregori L, Heger A, Jordan S, Behizad M, Reichl H, Romisch J, Svae T-  
 1226 E. Prion removal effect of a specific affinity ligand introduced into the manufacturing process of a  
 1227 pharmaceutical quality solvent/detergent (S/D)-treated plasma OctaplasLG®. *Vox Sanguinis*  
 1228 2009 2009;97:226-33.
- 1229 112. Heger A, Bailey A, Neisser-Svae A, Ertl M, Römisch J, Svae TE. Removal of prion infectivity by  
 1230 affinity ligand chromatography during OctaplasLG® manufacturing--results from animal bioassay  
 1231 studies. *Vox Sang.* 2012;102:294-301.
- 1232 113. Foster PR, Removal of TSE agents from blood products. *Vox Sang.* 2004 Jul;87 Suppl 2:7-10.
- 1233 114. Cai K, Gröner A, Dichtelmüller HO, Fabbrizzi F, Flechsig E, Gajardo R, von Hoegen I, Jorquera JI,  
 1234 Kempf C, Kreil TR, Lee DC, Moscardini M, Pölsler G, Roth NJ. Prion removal capacity of plasma



- 1235 protein manufacturing processes. A data collection from PPTA member companies. *Transfusion*.  
1236 2013;53:1894-905.
- 1237 115. Cardone F, Simoneau S, Arzel A, Puopolo M, Berardi VA, Abdel-Haq H, Galeno R, De Pascalis  
1238 A, Sbriccoli M, Graziano S, Valanzano A, Porte P, Diringer H, Brown P, Flan B, Pocchiari M.  
1239 Comparison of nanofiltration efficacy in reducing infectivity of centrifuged versus ultracentrifuged  
1240 263K scrapie-infected brain homogenates in "spiked" albumin solutions. *Transfusion*. 2012,  
1241 52:953-62.
- 1242 116. Seeger H, Heikenwalder M, Zeller N, Kranich J, Schwarz P, Gaspert A, Seifert B, Miele G, Aguzzi  
1243 A. Coincident scrapie infection and nephritis lead to urinary prion excretion. *Science* 2005;310:  
1244 324–326.
- 1245 117. Gregori L, Kovacs GG, Alexeeva I, Budka H, Rohwer RG. Excretion of transmissible spongiform  
1246 encephalopathy infectivity in urine. *Emerg Infect Dis*. 2008;14:1406-12.
- 1247 118. Haley NJ, Seelig DM, Zabel MD, Telling GC, Hoover EA. Detection of CWD Prions in Urine and  
1248 Saliva of Deer by Transgenic Mouse Bioassay. *PLoS ONE* 2009; 4(3): e4848.
- 1249 119. Gonzalez-Romero D, Barria MA, Leon P, Morales R, Soto C (2008) Detection of infectious prions  
1250 in urine. *FEBS Lett* 582: 3161–3166.
- 1251 120. Murayama Y, Yoshioka M, Okada H, Takata M, Yokoyama T, Mohri S. (2007) Urinary excretion  
1252 and blood level of prions in scrapie-infected hamsters. *J Gen Virol* 88: 2890–2898.
- 1253 121. Chen B, Morales R, Barria MA, Soto C. Estimating prion concentration in fluids and tissues by  
1254 quantitative PMCA. *Nat Methods*. 2010;7:519-20.
- 1255 122. Haley NJ, Mathiason CK, Carver S, Zabel M, Telling GC, Hoover EA. Detection of chronic wasting  
1256 disease prions in salivary, urinary, and intestinal tissues of deer: potential mechanisms of prion  
1257 shedding and transmission. *J Virol*. 2011;85:6309-18.
- 1258 123. Henderson DM, Denkers ND, Hoover CE, Garbino N, Mathiason CK, Hoover EA. Longitudinal  
1259 Detection of Prion Shedding in Saliva and Urine by Chronic Wasting Disease-Infected Deer by  
1260 Real-Time Quaking-Induced Conversion. *J Virol*. 2015;89:9338-47.
- 1261 124. John TR, Schätzl HM, Gilch S. Early detection of chronic wasting disease prions in urine of pre-  
1262 symptomatic deer by real-time quaking-induced conversion assay. *Prion*. 2013;7:253-8.
- 1263 125. Rubenstein R, Chang B, Gray P, Piltch M, Bulgin MS, Sorensen-Melson S, Miller MW. Prion  
1264 disease detection, PMCA kinetics, and IgG in urine from sheep naturally/experimentally infected  
1265 with scrapie and deer with preclinical/clinical chronic wasting disease. *J Virol*. 2011;85:9031-8.
- 1266 126. Notari S, Qing L, Pocchiari M, Dagdanova A, Hatcher K, Dogterom A, Groisman JF, Lumholtz IB,  
1267 Puopolo M, Lasmezas C, Chen SG, Kong Q, Gambetti P. Assessing prion infectivity of human  
1268 urine in sporadic Creutzfeldt-Jakob disease. *Emerg Infect Dis*. 2012;18:21-8.
- 1269 127. Luk C, Jones S, Thomas C, Fox NC, Mok TH, Mead S, Collinge J, Jackson GS. Diagnosing  
1270 Sporadic Creutzfeldt-Jakob Disease by the Detection of Abnormal Prion Protein in Patient  
1271 Urine. *JAMA Neurol*. 2016;73:1454-1460.
- 1272 128. Brandel JP, Galanaud D, Freeman L, Laplanche JL, Haik S Variant or sporadic Creutzfeldt-Jakob  
1273 disease? *The Lancet*, Vol. 375 No. 9718 p 889; 2010.
- 1274 129. AFSSaPS

- 1275 129a Analyse du risque de transmission de la variante de la maladie de Creutzfeldt-Jakob  
1276 (vMCJ) et de la forme sporadique de la maladie de Creutzfeldt-Jakob par les produits de santé  
1277 d'origine humaine– Septième actualisation des travaux du groupe d'experts pluridisciplinaire.  
1278 Rapport de juillet 2009. [http://www.afssaps.fr/Dossiers-thematiques/Creutzfeldt-Jakob-et-  
produits-de-sante](http://www.afssaps.fr/Dossiers-thematiques/Creutzfeldt-Jakob-et-<br/>1279 produits-de-sante)
- 1280 129b Analyse du risque de transmission de la variante de la maladie de Creutzfeldt-Jakob (vMCJ)  
1281 et de la forme sporadique de la maladie de Creutzfeldt-Jakob par les produits de santé d'origine  
1282 humaine– Sixième actualisation des données du rapport du groupe d'experts adhoc de décembre  
1283 2000 – Rapport de novembre 2007. [http://www.afssaps.fr/Dossiers-thematiques/Creutzfeldt-  
Jakob-et-produits-de-sante](http://www.afssaps.fr/Dossiers-thematiques/Creutzfeldt-<br/>1284 Jakob-et-produits-de-sante)
- 1285 129c Evaluation of the risk of transmission of Creutzfeldt-Jakob disease agent by blood and its  
1286 constituents – Experts' group meeting of 16 november 2004. [http://www.afssaps.fr/Dossiers-  
thematiques/Creutzfeldt-Jakob-et-produits-de-sante](http://www.afssaps.fr/Dossiers-<br/>1287 thematiques/Creutzfeldt-Jakob-et-produits-de-sante)
- 1288 129d Analysis of the risk of transmission of variant of Creutzfeldt-Jakob disease by health  
1289 products and by tissues and fluids of human origin – Update of findings of ad hoc group report of  
1290 December 2000 – Report of February 2004. [http://www.afssaps.fr/Dossiers-  
thematiques/Creutzfeldt-Jakob-et-produits-de-sante](http://www.afssaps.fr/Dossiers-<br/>1291 thematiques/Creutzfeldt-Jakob-et-produits-de-sante)
- 1292 129e Analysis of the risk of transmission of the variant Creutzfeldt-Jakob disease by medicinal  
1293 products products of human origin and labile blood products – Data update of the ad hoc group  
1294 report dated December 2000 –March 2003 report. [http://www.afssaps.fr/Dossiers-  
thematiques/Creutzfeldt-Jakob-et-produits-de-sante](http://www.afssaps.fr/Dossiers-<br/>1295 thematiques/Creutzfeldt-Jakob-et-produits-de-sante)
- 1296 129f Analysis of the risk of transmission of the variant Creutzfeldt-Jakob disease by medicinal  
1297 products products of human origin and labile blood products – Data update of the ad hoc group  
1298 report dated December 2000 –February 2002 report. [http://www.afssaps.fr/Dossiers-  
thematiques/Creutzfeldt-Jakob-et-produits-de-sante](http://www.afssaps.fr/Dossiers-<br/>1299 thematiques/Creutzfeldt-Jakob-et-produits-de-sante)
- 1300 129g Analysis of the risk of transmission of new variant Creutzfeldt-Jakob disease by blood and  
1301 its derivatives”, Recommendations of expert group convened at the initiative of AFSSaPS, 11  
1302 December 2000. <http://afssaps.sante.fr>
- 1303 129h Revision of measures to minimise the risk of TSE transmission via blood products”, Report  
1304 of expert group convened under the aegis of the Agence Française de Sécurité Sanitaire des  
1305 Produits de Santé (AFSSaPS) and the Etablissement Français du Sang (EFS), February 2000.  
1306 <http://afssaps.sante.fr>