

MW/SAN

12th December 1979.

Dr. M. Chisholm,  
Consultant Haematologist,  
Royal South Hants Hospital,  
SOUTHAMPTON

Dear Dr. Chisholm,

re: Gary WEBSTER GRO-A 65

GRO-A

I enclose a copy of Gary's bleeding episodes so far this term. He bleeds mainly into his knees and elbows.

He was seen by Mr. Moynihan on 20.9.79 to reassess his left knee following traction over the summer holiday. Mr. Moynihan advised he should wear his block leather splint during the day in between bleeds for three to six months and to wear it at night when he bleeds into his left knee.

Gary used to have full range of movement despite his deformed knees, but now he seems to have lost some range of movement due to "deterioration arthropathy" during the past few years. The range of movement of his right and left knees are - 10° - 150° and - 15° - 150° respectively.

This term he has had an intensive programme of physiotherapy for both quads, but was unfortunately interrupted by the frequent bleeds into his knees. He has not yet recovered his full range of movement of the left knee.

Biochemically, his S.G.O.T. has been persistently raised for the past nine months ranging from just about the upper limit of normal range to 62 IU/L.

He is now competent in self therapy.

Yours sincerely,

M. Wassef  
Senior Clinical Medical Officer.

ENC.