

Table 1**Glasgow Yorkhill Haemophilia Centre****Comparison of product use, 1979 - 1987 (excluding Factor IX and FFP)**

| Year | Cryoprecipitate | | F8 concentrate | | FEIBA and DDAVP | | Porcine F8 | | Total Units |
|------|-----------------|-------|----------------|--------|---------------------|---|--------------|------|-------------|
| | <i>units</i> | % | <i>units</i> | % | <i>units</i> | % | <i>units</i> | % | |
| 1979 | 60 | 0.01% | 491,741 | 99.99% | Nil | | Nil | | 491,801 |
| 1980 | 1,305 | 0.1% | 843,974 | 99.9% | Nil | | Nil | | 845,279 |
| 1981 | 29,200 | 3% | 1,089,623 | 97% | Nil | | Nil | | 1,118,823 |
| 1982 | 7,050 | 0.7% | 1,002,180 | 99.3% | Nil | | Nil | | 1,009,230 |
| 1983 | 30,500 | 2.5% | 1,157,925 | 97.5% | Nil | | Nil | | 1,188,425 |
| 1984 | 32,340 | 3% | 1,040,856 | 97% | Units not specified | | Nil | | 1,073,196 |
| 1985 | 27,930 | 3.6% | 739,094 | 96.4% | Nil | | Nil | | 767,024 |
| 1986 | 38,780 | 6% | 600,360 | 93.5% | Nil | | 2,720 | 0.5% | 641,860 |
| 1987 | 27,860 | 3% | 922,700 | 97% | Units not specified | | Nil | | 950,560 |

Table 2**Glasgow Yorkhill Haemophilia Centre****Source of F8 concentrate**

| Year | PFC <i>units</i> | PFC % | Commercial <i>units</i> | Commercial % | Total units |
|------|---------------------|----------|----------------------------|-----------------|-------------|
| 1979 | 137,465 | 28% | 354,276 | 72% | 491,741 |
| 1980 | 161,242 | 19% | 682,732 | 81% | 843,974 |
| 1981 | 457,126 | 42% | 632,497 | 58% | 1,089,623 |
| 1982 | 516,300 | 51.5% | 485,880 | 48.5% | 1,002,180 |

| | | | | | |
|-------------|-----------|-------|--------|------|-----------|
| 1983 | 1,121,075 | 97% | 36,850 | 3% | 1,157,925 |
| 1984 | 1,035,396 | 99.5% | 5,460 | 0.5% | 1,040,856 |
| 1985 | 739,094 | 100% | Nil | 0% | 739,094 |
| 1986 | 600,360 | 100% | Nil | 0% | 600,360 |
| 1987 | 922,700 | 100% | Nil | 0% | 922,700 |

Source: National Haemophilia Database Bleeding Disorder Statistics for the Penrose Inquiry
[PRSE0002887]