

**CL Table 1****Edinburgh Haemophilia Centre****Comparison of product use, 1979 - 1984 (excluding Factor IX and FFP)**

| Year | Cryoprecipitate |     | F8 concentrate |     | FEIBA and DDAVP |    | Porcine F8   |    |
|------|-----------------|-----|----------------|-----|-----------------|----|--------------|----|
|      | <i>units</i>    | %   | <i>units</i>   | %   | <i>units</i>    | %  | <i>units</i> | %  |
| 1979 | 694,190         | 78% | 201,486        | 22% |                 |    |              |    |
| 1980 | 1,212,470       | 40% | 1,808,750      | 60% |                 |    |              |    |
| 1981 | 720,160         | 36% | 1,282,148      | 64% |                 |    |              |    |
| 1982 | 612,000         | 26% | 1,620,100      | 69% | 110,000         | 5% |              |    |
| 1983 | 341,700         | 15% | 1,907,000      | 83% | 37,000          | 2% |              |    |
| 1984 | 139,000         | 5%  | 2,545,380      | 91% | 8,006           | 0% | 91,200       | 3% |

**CL Table 2****Edinburgh Haemophilia Centre****Source of F8 concentrate**

| Year | PFC          | PFC    | Commercial   | Commercial |
|------|--------------|--------|--------------|------------|
|      | <i>units</i> | %      | <i>units</i> | %          |
| 1979 | 201,486      | 100.0% | 0            | 0.0%       |
| 1980 | 1,644,750    | 90.9%  | 164,000      | 9.1%       |
| 1981 | 840,130      | 65.5%  | 442,018      | 34.5%      |
| 1982 | 1,612,800    | 99.5%  | 7,300        | 0.5%       |
| 1983 | 1,756,000    | 92.1%  | 151,000      | 7.9%       |
| 1984 | 2,506,880    | 98.5%  | 38,500       | 1.5%       |

Source: National Haemophilia Database Bleeding Disorder Statistics for the Penrose Inquiry  
[PRSE0002887]