

Date

Left knee

Treatments 1974 (donor units cryoprecipitate)

19.2.74	5	Right ankle
19.2.74	10 (aspirated)	11.3.74 5
20.2.74	8	12.3.74 6
21.2.74	5	
25.2.74	5	
28.2.74	5	
22.4.74	5	
24.4.74	5	
3.9.74	8	
24.9.74	8	
2.10.74	8	
15.10.74	600 Lister	
21.10.74	720 "	
22.10.74	720 "	
28.10.74	400 "	
29.10.74	8 Cryo	
1.11.74	800 Lister	
4.11.74	6 Cryo	
5.11.74	6	
6.11.74	6	
7.11.74	5	
8.11.74	6	
10.11.74	6	
11.11.74	6	
12.11.74	6	
13.11.74	6	
14.11.74	5	
15.11.74	5	
18.11.74	5	
19.11.74	5	
20.11.74	5	
21.11.74	5	
22.11.74	5	
25.11.74	5	
26.11.74	5	
27.11.74	5	
28.11.74	5	

FIRST RECORD OF FVIII

GRO-C: D A Pollard

Treatments 1975 (donor units cryoprecipitate)

19.3.75	15 aspirated	Left elbow	Right elbow
20.3.75	8	13.4.75	6.1.75 8
21.3.75	8		10.2.75 8
22.3.75	6		12.2.75 8
25.3.75	6		
26.3.75	5		
1.4.75	6		
2.4.75	6		
16.4.75	12		
17.4.75	10		
18.4.75	10		

Treatments 1976 (donor units cryoprecipitate?)

21.4.76	500 Lister	Right ankle	Right shoulder
22.4.76	880 "	1.4.76 560 units	14.3.76 566 units

Finger

30.4.76 (cut) 530 Liser

Left thigh?

7.6.76 265 units Lister

PLEASE RECORD EVERY VISIT

WEIGHT IN KG.

(3)

NAME		GRO-B		HOSPITAL NO.		DATE WEIGHED											
DATE	TIME	IP OR OP	CLINICAL NOTES Time of onset, severity splints, crutches etc.)	GIVEN BY	DRUGS EACA Cyclokapron Antibiotics	TYPE Cryo:FFP IX conc. VIII " Blood etc	GROUP (if cryo or FFP)	DONOR NO BATCH NO	VOL	PRE %	POST %	DOSE %	DOSE U.	U Kg	EX RISE	RE- COVER	
4.5.76	11.30	OP	Swollen onset of CORRINGHAM pain (D) tip of shoulder blade after swollen movement over Tender, post used. aspect scapula pain related to movement involving (C) trapezius ? trapezius bleed. (C)			VIII Conc —		190875 190875	36 35	<div style="border: 1px solid black; padding: 5px; display: inline-block;">GRO-C</div>							