

**The Liver Unit**  
**Royal Victoria Hospital**  
**1<sup>st</sup> Floor**  
**East Wing**  
**Grosvenor Road**  
**Belfast**  
**BT12 6BA**

**Consultants**  
**Dr Neil McDougall, Consultant Gastroenterologist/Hepatologist**  
**Dr Ian Cadden, Consultant Hepatologist**  
**Dr Jonathan Cash, Consultant Hepatologist**  
**Dr Roger McCorry, Consultant Hepatologist**  
**Dr Conor Braniff, Consultant Hepatologist**  
**Tel: 02890 [redacted] Dr McDougall's Secretary**  
**02890 [redacted] Dr Cadden's Secretary**  
**02890 [redacted] Dr Cash's Secretary**  
**02890 [redacted] Dr McCorry's Secretary**  
**02890 [redacted] Dr C Braniff's Secretary**

**Miss Mary Bright**  
**Mrs Teresa Gault**  
**Mrs Siobhan Miskelly**  
**Mrs Emma Chapman**  
**Ms Rosemary Smyth**

Hospital No: RV.84/004361  
H&C No.: **GRO-A**

Date Typed: 17/01/2019  
Date Dictated: 14/01/2019

DR LWJ MCNEILL  
CITYVIEW MEDICAL  
WATERSIDE HEALTH CENTRE  
127 SPENCER ROAD  
LONDONDERRY  
BT47 6AH

Dear DR MCNEILL

**RE: MR. EDWARD FRANCIS CONWAY, D.O.B. [redacted] GRO-C1958**  
**GRO-C**

Thank you for your letter dated 20 December 2018 and for your efforts to try and reassure this gentleman that he does not have evidence of liver cancer at present. I believe our Trust will be seeking to arrange a meeting with Mr Conway to try and clarify his long-term follow-up arrangements and I have offered to take part in that meeting.

In answer to your specific questions:

1. I would have no objection to you starting him on a statin despite the diagnosis of probable liver cirrhosis. We often use statins in patients with liver cirrhosis. The only condition would be that it would be wise to monitor his liver tests every couple of weeks for the first six weeks or so to ensure that the statin does not cause any upset of transaminases.
2. With respect to follow-up imaging, the UK National Guidelines currently state that screening for hepatocellular cancer should be offered with ultrasound every 6 months and not anymore frequently than that (for patients with risk factors such as liver cirrhosis). Therefore he should have a repeat ultrasound roughly 6 months after his last imaging and not any sooner. I am afraid we cannot do more frequent scanning just because of a relative's request.
3. Regarding long-term follow-up, I will be happy to facilitate this through the Liver Unit in the Royal if the patient and his family are willing to comply with appointments. I appreciate that this is slightly contradictory given that his sister stated that they do not have any faith or confidence in the tests that we arrange or the investigations that we carry out. Hopefully we will be able to resolve this matter by arranging a meeting with the family.

Mr Conway should currently be on medication to treat his chronic hepatitis C. I will liaise with our Nurse Specialists to clarify the date that he finishes his hepatitis C treatment. He will require a follow-up blood test 12 weeks after completion of treatment to help clarify whether or not the virus has been successfully cleared. I will arrange his next review at our clinic with the result of his follow-up blood test so we can discuss the result. His follow-up ultrasound should be around the same time.

Kind regards

Yours sincerely

Dr Neil McDougall MD FRCP (Ed)  
Consultant Gastroenterologist/Hepatologist

CC Karen Patterson/ Orla McCormick Specialist Hepatology Nurses Ambulatory Care Centre RVH

/MGB

MR EDWARD FRANCIS CONWAY H&C: 606 013 0801