

GRO-A

Haemophiliac with inhibitors.

Bottle No.	Infusion	Additions	Score	Final fluids
DATE	CLINICAL INDICATION AND/OR TREATMENT		RESULTS	
2.2.73	Not responding clinically to Cryoprecipitate. Trial dose of Cryoprecipitate given.		Factor VIII Pre-infusion: 0% Factor VIII Post-infusion: 0% Factor VIII Inhibitor Titre: 1/10	
4.2.73			Factor VIII Assay: 0%	
6.3.73	Trial dose of Haemofil given.		Factor VIII Assay: 30%	
7.3.73			Factor VIII Assay: 8%	
9.3.73 - 10.3.73	Given Cryoprecipitate.			
18.3.73	Treatment reverted to Haemofil.			
19.3.73			Factor VIII Assay: 8%	
			Factor VIII Inhibitor Titre: 1/20	
20.3.73			Factor VIII Assay: 5.5%	
			No inhibitors detected against Haemofil.	
25.5.73			Factor VIII Assay: 0%	
			Inhibitor titre against <u>Cryoprecipitate</u> : 1/20	
2.7.73	(Specimen taken 28.6.73). After approx. one month's continuous treatment on Haemofil.		No inhibitors detected against Haemofil.	
11.1.74			(With Haemofil) Pre-infusion Factor VIII Assay: 7% Post-infusion Factor VIII Assay: 22%	
12.1.74			Factor VIII Assay: 14%	
29.1.74	Given double dose of Haemofil		Pre-infusion Factor VIII Assay: 0% Post-infusion Factor VIII Assay: 70%	
7.2.74			Post-infusion Factor VIII Assay: 32.5%	
28.2.74	12 hours post-infusion with Haemofil		Factor VIII Assay: 30%	