

29<sup>th</sup> ANNUAL REPORT 2020

# **CREUTZFELDT-JAKOB DISEASE SURVEILLANCE IN THE UK**

---

**The National CJD Research & Surveillance Unit**

Western General Hospital, Edinburgh, EH4 2XU  
&  
Chancellor's Building, Royal Infirmary of Edinburgh, EH16 4SB

[www.cjd.ed.ac.uk](http://www.cjd.ed.ac.uk)

# Table of Contents

## SECTION 1

Summary	3
---------	---

## SECTION 2

Clinical Surveillance	5
2.1 Referrals	5
2.2 Sporadic CJD	6
2.3 Variant CJD	12
2.4 Iatrogenic CJD	16
2.5 Transfusion Medicine Epidemiology Review (TMER)	16
2.6 Study of Progressive Intellectual and Neurological Deterioration (PIND)	17
2.7 Surveillance of potential occupational exposure to CJD	18
2.8 Prion surveillance in primary immunodeficiency patients	18
2.9 Enhanced surveillance of those at increased risk of CJD	18

## SECTION 3

Laboratory Activities	20
3.1 Neuropathology – Statement of Progress and Surveillance Activities	20
3.2 Protein Biochemistry Laboratory	22
3.3 Brain Banking Activities	23
3.4 Molecular Genetics	24
3.5 CSF RT-QuIC, 14-3-3 and other brain specific proteins	25

## SECTION 4

National Care Team	27
--------------------	----

## SECTION 5

Publications	29
--------------	----

## SECTION 6

Staff	31
-------	----

## SUMMARY

**T**he national surveillance programme for Creutzfeldt-Jakob disease (CJD) in the UK was initiated in May 1990. In 1999, the National CJD Research & Surveillance Unit (NCJDRSU) became a WHO Collaborative Centre on the surveillance, diagnosis and epidemiology of human transmissible spongiform encephalopathies (TSEs). In September 2001, the National Care Team was formed in response to concerns regarding the care of CJD patients. The team currently comprises two care coordinators (who are senior nurses) with secretarial and clinical neurological support from within the NCJDRSU where it is based.

The annual mortality rate for sporadic CJD (sCJD) was 1.98 cases/million in 2020. Although the data for 2020 may still be incomplete, detailed clinical and epidemiological information has been obtained for the great majority of patients. Although the autopsy rate in cases of suspected CJD has decreased in recent years, it remains relatively high in comparison to the general autopsy rate in the UK. The number of brain tissue specimens examined for sCJD in the neuropathology laboratory in 2020 was 12 cases (compared with 15 cases in 2019).

Over the period 1990-2020 average annual mortality rates from sCJD in England, Wales, Scotland and Northern Ireland were, respectively, 1.24, 1.51, 1.24 and 0.95/million/year. The differences between these rates are not statistically significant ( $p=0.41$ ). The mortality rates of sCJD in the UK are comparable to those observed in most other European countries and elsewhere in the world, including countries that are free of BSE.

Variably Protease Sensitive Prionopathy (VPSPr), is of uncertain nosological significance but is presently considered a form of sporadic prion disease, alongside sCJD. The NCJDRSU has so far identified a total of 20 such cases in the UK and is continuing to monitor this form of disease.

Up to 31<sup>st</sup> December 2020, 178 cases of definite or probable variant CJD (vCJD) had been identified in the UK (123 definite and 55 probable cases who did not undergo post mortem). All 178 cases have died. The clinical, neuropathological and epidemiological features of the cases of vCJD are remarkably uniform and consistent with previous descriptions. Risk factors for the development of vCJD include age, residence in the UK and methionine homozygosity at Codon 129 of the prion protein gene – of 161 clinically affected definite and probable cases of vCJD with available genetic analysis, 160 have been methionine homozygotes and one methionine-valine heterozygous at Codon 129 of the *PRNP* gene. Analysis of vCJD diagnoses and deaths from January 1994 continues to indicate that the peak has passed. While this is an encouraging finding, the incidence of vCJD may increase again, particularly if further cases in different genetic subgroups with longer incubation periods exist. The identification of an individual of the *PRNP*-129 MV genotype as a confirmed case of vCJD (in addition to the possible case of vCJD reported in the NCJDRSU 17<sup>th</sup> Annual Report, 2008) and the finding of disease-related prion protein in the spleen of a clinically unaffected blood recipient (reported in 2004) is consistent with such a hypothesis. These cases, along with the results of large-

scale surveys of the prevalence of abnormal prion protein in appendix and tonsil tissues suggest the possibility of a greater number of asymptomatic infections (either preclinical or subclinical) in the population than might be indicated by the present numbers of confirmed clinical cases.

To help prevent any possible spread of CJD between people, we continue to ask clinicians to refer all new suspect CJD cases to their local infection control and health protection teams. This is important as a local response may be required with respect to limiting potential secondary transmission and other issues that may arise over time concerning the protection of the wider community. The NCJDRSU continues to assist local health protection teams in local audit and investigations of cases in response to local concerns. The NCJDRSU also continues to collaborate with government health departments and the UK public health authorities, including Public Health England and Health Protection Scotland, in a range of activities in relation to the follow up of those identified as at increased risk of CJD.

The activities of the NCJDRSU are strengthened by collaboration with other surveillance projects, including the Transfusion Medicine Epidemiology Review, Prion Surveillance in Primary Immunodeficiency Patients, the study of Progressive Intellectual and Neurological Deterioration in Children and the CJD International Surveillance Network. The collaboration of our colleagues in these projects is greatly appreciated; the effectiveness of this collaboration allowed the identification in 2003 of a case of vCJD associated with blood transfusion and the identification in 2004 of disease-related PrP in the spleen of a recipient of blood donated by someone incubating vCJD. A patient was also identified in 2010 who had evidence of vCJD infection in the spleen (but no evidence of clinical vCJD), considered probably due to blood products (treatment for haemophilia).

The data concerning CSF RT-QuIC are given in Section 3.5; the sensitivity of CSF RT-QuIC for a diagnosis of sCJD is comparable with that of CSF 14-3-3. The specificity is superior to that of CSF 14-3-3 with no positives in cases with a confirmed alternate diagnosis.

The success of the National CJD Research & Surveillance Unit continues to depend on the extraordinary level of co-operation from the neurology and neuropathology communities and other medical and paramedical staff throughout the UK. Ongoing support is provided by the Infectious Disease Epidemiology Unit, London School of Hygiene and Tropical Medicine. We are also particularly grateful to the relatives of patients for their collaboration.

Providing information to the public is an important aspect of the NCJDRSU's activities. We liaise closely with the CJD Support Network, providing articles for their newsletter, updating their information booklets and giving presentations to their Annual Family Day meetings. Professor Knight is the current Chair of the Network's Management Committee and is also a member of the CJD International Alliance of CJD support organisations.




## CLINICAL SURVEILLANCE

The national surveillance of CJD in the UK was initiated in May 1990. Surveillance is funded by the Department of Health and Social Care, UK and by the Scottish Government Health Department. The NCJDRSU aims to monitor characteristics of CJD, specifically sCJD and vCJD, to identify trends in incidence rates and to study risk factors for the development of disease. This report documents the findings in relation to UK cases of sCJD and vCJD as well as genetic and iatrogenic forms of disease referred up to 31st December 2020 (based on data ascertained up to 3<sup>rd</sup> July 2021). Mortality data from England and Wales include retrospective data from 1970; for Scotland and Northern Ireland, retrospective mortality data are available from 1985. Case definitions for the various types of CJD can be found at [www.cjd.ed.ac.uk/sites/default/files/criteria.pdf](http://www.cjd.ed.ac.uk/sites/default/files/criteria.pdf). Cases classified as definite or probable are included in all analyses from Section 2.2 onwards.

### 2.1 Referrals to NCJDRSU

The NCJDRSU receives referrals of suspect cases of CJD and a proportion of these will turn out not to have CJD. Referrals of suspect cases increased after the present surveillance system began in 1990, particularly following the description of vCJD in 1996. Numbers of referrals fluctuate over time, and may be attributed to variation in case ascertainment and reporting practice, including changes in the number of non-CJD cases referred to the NCJDRSU (see Figure 1)


**Figure 1** Referrals to NCJDRSU: UK 1 May 1990 – 31<sup>st</sup> December 2020



In addition to formal referrals of suspected CJD as shown above, the NCJDRSU also receives enquiries from clinicians for advice or to utilise the CSF tests available at the Unit. In 2020, in addition to the 168 formal referrals of suspected cases of CJD, there were a further 238 enquiries where advice was sought by clinicians on individual patients. The NCJDRSU also receives a number of enquiries from relatives of patients, members of the public and professional bodies both in the UK and worldwide seeking advice in relation to CJD. In 2020, over 110 such email enquiries were received via the contact details displayed on our website (<http://www.cjd.ed.ac.uk/contact-us>).

Nationally across the UK rates of post-mortem examinations have been decreasing over time, and this includes for suspected cases of CJD (Figure 2). Although increasing diagnostic certainty can now be offered by biomarker tests (MRI, RT-QuIC), the fall in post-mortem rate may potentially impact on our ability to confirm the different types of prion disease, particularly in cases where prion disease may not have been considered or if otherwise atypical of CJD.

**Figure 2** Post-mortem rate in all referrals of suspected CJD to NCJDRSU: UK  
1 May 1990 – 31<sup>st</sup> December 2020



## 2.2 Sporadic Creutzfeldt-Jakob Disease

Between 1st January 1970 and 31st December 2020, 2762 cases of sCJD were identified (268 in England and Wales from 1970-1984 and 2494 in the UK from 1985-2020), of which 26 cases were alive on 31st December 2020. Three cases moved abroad after diagnosis and are therefore lost to follow-up. Of these 2762 cases, 1624 (59%) were classified as definite cases with the remainder classed as probable; 1396 (51%) were female and 1366 (49%) were male. Eight further cases have been identified: 3 in Jersey, 3 in the Isle of Man and 2 cases who were repatriated to the UK when they became ill but had been living abroad. These 8 cases are not included in the following UK analyses.

Figure 3 shows the annual mortality rates from sCJD for the UK between 1985 and 2020. The number of deaths identified each year has increased over time. A similar phenomenon has been observed in other European countries, and may reflect improved case ascertainment, particularly in those aged over 70 years, and also following revised diagnostic criteria in January 2017, the impact of which is under investigation.

**Figure 3 Mortality Rates from sCJD, UK, 1985-2020**





Figure 4 shows average annual age-specific mortality rates over the time periods 1970-1984, 1985-1994, 1995-2004, 2005-2014 and subsequently. These data also emphasise the very small numbers of cases of sCJD occurring in individuals aged <50 years. The median ages of cases at death during these five time periods were 63, 65, 67, 69 and 69½ years, respectively. In all five time periods, the mortality rates below 40 years of age were low (< 0.04/million/year). Thereafter, in all five periods, the mortality rates increased up to ages 65-79 years and then declined. The reasons for this decline are unclear but might be explained in part by an under-ascertainment in the most elderly. Comparison between the different time periods, indicate an increase in age-specific recorded mortality over time in all age groups over 50. These observations are consistent with improved case ascertainment in all ages over 50 years, but with the greatest increase occurring in the elderly.

**Figure 4** Age-specific mortality rates from sporadic CJD in the UK 1970-2020  
(note: from 1970-1984 only England and Wales, thereafter UK)



1970-1984 Mortality rates calculated using mid-1981 England & Wales population estimates based on the 1981 Census  
 1985-1994 Mortality rates calculated using mid-1991 UK population estimates based on the 1991 Census  
 1995-2004 Mortality rates calculated using mid-2001 UK population estimates based on the 2001 Census  
 2005-2014 Mortality rates calculated using mid-2011 UK population estimates based on the 2011 Census  
 2015-2020 Mortality rates calculated using mid-2011 UK population estimates based on the 2011 Census


## Geographical distribution of sCJD

Over the period 1990-2020 the average crude annual mortality rates from sCJD per million population were 1.24 in England, 1.51 in Wales, 1.24 in Scotland and 0.95 in Northern Ireland (Tables 1a and 1b). When account is taken of age and sex, the variation in recorded mortality between the different countries is not statistically significant ( $p=0.41$ ).

Age- and sex- standardised mortality ratios (SMRs) for the 12 government office regions of the UK for the period 1st January 1990 to 31st December 2020 were calculated (Figure 5). An SMR of 100 equates to the national average mortality rate; an SMR above or below this value reflects relative high or low mortality, respectively. After adjusting for the age/sex distribution of the population, the variation in mortality rates between the different regions is not statistically significant ( $p=0.14$ ).



**Figure 5**      **Standardised sporadic CJD mortality ratios (SMRs)**  
1 January 1990 - 31 December 2020, by region of residence at death



**Table 1a Deaths from definite and probable sporadic CJD in England (shown by region and local authority of residence at death). 1<sup>st</sup> January 1990 to 31<sup>st</sup> December 2020**

ENGLAND	No. of cases	Mortality Rate*	ENGLAND	No. of cases	Mortality Rate*
<b>North East</b>	<b>90</b>	<b>1.14</b>	<b>East</b>	<b>235</b>	<b>1.36</b>
Darlington UA	5		Luton UA	3	
Hartlepool UA	3		Peterborough UA	4	
Middlesbrough UA	1		Southend-on-Sea UA	5	
Redcar & Cleveland UA	6		Thurrock UA	4	
Stockton-on-Tees UA	5		Bedfordshire	18	
Durham	15		Cambridgeshire	14	
Northumberland	14		Essex	75	
Tyne & Wear	41		Hertfordshire	38	
			Norfolk	39	
<b>North West</b>	<b>265</b>	<b>1.24</b>	Suffolk	35	
Blackburn with Darwen UA	8		<b>London</b>	<b>220</b>	<b>0.94</b>
Blackpool UA	3		Inner London	66	
Halton UA	7		Outer London	154	
Warrington UA	11		<b>South East</b>	<b>342</b>	<b>1.34</b>
Cheshire	24		Bracknell Forest UA	3	
Cumbria	22		Brighton and Hove UA	2	
Greater Manchester	83		Isle of Wight UA	4	
Lancashire	48		Medway UA	7	
Merseyside	59		Milton Keynes UA	4	
<b>Yorkshire and the Humber</b>	<b>182</b>	<b>1.15</b>	Portsmouth UA	4	
East Riding of Yorkshire UA	12		Reading UA	7	
Kingston Upon Hull, City of UA	5		Slough UA	1	
North East Lincolnshire UA	6		Southampton UA	5	
North Lincolnshire UA	6		West Berkshire UA	8	
York UA	8		Windsor and Maidenhead UA	5	
North Yorkshire	31		Wokingham UA	6	
South Yorkshire	50		Buckinghamshire	14	
West Yorkshire	64		East Sussex	26	
<b>East Midlands</b>	<b>163</b>	<b>1.21</b>	Hampshire	58	
Derby UA	13		Kent	73	
Leicester UA	12		Oxfordshire	33	
Nottingham UA	8		Surrey	38	
Rutland UA	2		West Sussex	44	
Derbyshire	32		<b>South West</b>	<b>261</b>	<b>1.66</b>
Leicestershire	24		Bath & North East Somerset UA	8	
Lincolnshire	27		Bournemouth UA	8	
Northamptonshire	14		Bristol, City of UA	12	
Nottinghamshire	31		North Somerset UA	15	
<b>West Midlands</b>	<b>195</b>	<b>1.17</b>	Plymouth UA	18	
Herefordshire, County of UA	7		Poole UA	6	
Stoke-on-Trent UA	6		South Gloucestershire UA	14	
Telford and Wrekin UA	6		Swindon UA	3	
Shropshire	11		Torbay UA	4	
Staffordshire	42		Cornwall and Isles of Scilly	32	
Warwickshire	11		Devon	34	
West Midlands (Met County)	87		Dorset	23	
Worcestershire	25		Gloucestershire	30	
			Somerset	32	
			Wiltshire	22	
<b>TOTAL FOR ENGLAND</b>	<b>1953</b>	<b>1.24</b>			

\* number of deaths/million/annum based on mid-2005 population estimates in England (source: ONS) over the 31-year period of the study. Postcode of residence obtained from AFD Postcode Plus.

**Table 1b Deaths from definite and probable sporadic CJD: Wales, Scotland and NI**  
**1<sup>st</sup> January 1990 to 31<sup>st</sup> December 2020**

<b>WALES†</b>	<b>No. of cases</b>	<b>WALES†</b>	<b>No. of cases</b>
Isle of Anglesey	5	Neath Port Talbot	2
Gwynedd	7	The Vale of Glamorgan	4
Conwy	6	Cardiff	11
Denbighshire	2	Bridgend	3
Flintshire	6	Rhondda, Cynon, Taff	9
Wrexham	8	Merthyr Tydfil	3
Powys	16	Caerphilly	13
Ceredigion	3	Blaenau Gwent	1
Pembrokeshire	3	Torfaen	6
Cardiganshire	6	Monmouthshire	4
Swansea	14	Newport	7
<b>TOTAL FOR WALES</b> <b>(MORTALITY RATE*)</b>	<b>139</b> <b>(1.51)</b>	†unitary authorities	

<b>SCOTLAND†</b>	<b>No. of cases</b>	<b>SCOTLAND†</b>	<b>No. of cases</b>
Aberdeen City	7	Highland	6
Aberdeenshire	12	Inverclyde	3
Angus	3	Midlothian	2
Argyll & Bute	4	Moray	5
Clackmannanshire	2	North Ayrshire	4
Dumfries & Galloway	6	North Lanarkshire	9
Dundee City	9	Orkney Islands	0
East Ayrshire	3	Perth & Kinross	4
East Dunbartonshire	4	Renfrewshire	6
East Lothian	4	Scottish Borders	4
East Renfrewshire	3	Shetland Islands	3
Edinburgh, City of	27	South Ayrshire	5
Eilean Siar	1	South Lanarkshire	10
Falkirk	5	Stirling	7
Fife	11	West Dunbartonshire	5
Glasgow City	18	West Lothian	5
<b>TOTAL FOR SCOTLAND</b> <b>(MORTALITY RATE*)</b>	<b>197</b> <b>(1.24)</b>	†council areas	


<b>NORTHERN IRELAND†</b>	<b>No. of cases</b>	<b>NORTHERN IRELAND†</b>	<b>No. of cases</b>
Antrim and Newtonabbey	3	Fermanagh and Omagh	3
Ards and North Down	1	Lisburn and Castlereagh	5
Armagh City, Banbridge & Craigavon	8	Mid and East Antrim	2
Belfast	8	Mid Ulster	4
Causeway Coast and Glens	6	Newry, Mourne and Down	4
Derry City and Strabane	7		
<b>TOTAL FOR N IRELAND</b> <b>(MORTALITY RATE*)</b>	<b>51</b> <b>(0.95)</b>	†local government districts	

\* based on mid-2005 population estimates for unitary authorities in Wales, council areas within Scotland and local government districts in Northern Ireland (ONS) over the 31-year period of the study. Postcode of residence obtained from AFD Postcode Plus.

## 2.3 Variant Creutzfeldt-Jakob Disease

Up to 31<sup>st</sup> December 2020, 178 cases of definite or probable vCJD had been identified in the UK (123 definite and 55 probable cases who did not undergo post mortem). Seventy-five (42%) of the 178 cases were female and 103 (58%) were male. The median age at onset of disease was 26½ years and the median age at death 28 years (compared with 67 years for the median age at onset and 68 years for the median age at death for sCJD). The youngest case was aged 12 years at onset while the oldest case was aged 74 years. The age- and sex-specific mortality rates for vCJD over the time period 1 May 1995 to 31 December 2020 are shown in Figure 6. The median duration of illness from the onset of first symptoms to death was 14 months (range 6-114) compared with a median duration of illness for cases of sCJD of 4 months (range 1 to 132) during the period 1990-2019. The last known UK case of vCJD was reported in 2016 with onset in 2014.

**Figure 6** Age- and sex-specific mortality rates from variant CJD in the UK  
1 May 1995 - 31st December 2020



Mortality rates calculated using ONS mid-2001 population estimates

Of 161 vCJD cases tested, one case of definite vCJD was heterozygous (MV) at Codon 129 of the *PRNP* gene while the remaining 160 definite or probable vCJD cases were methionine homozygous (MM). A single case of possible vCJD with an MV genotype was described by Kaski et al. in 2009.<sup>1</sup> To date, no case of vCJD has been identified in the UK in individuals born after 1989.

### Geographical distribution of variant CJD

Tables 2a and 2b present data on the geographical distribution by residence at onset (for all 178 vCJD cases) and residence at death (for 175 vCJD cases who had died and were resident in the UK at death), along with the crude mortality rate per million population per annum of each standard region.

<sup>1</sup> Kaski D, Mead S, Hyare H, Cooper S, Jampana R, Overell J, Knight R, Collinge J, Rudge P: Variant CJD in an individual heterozygous for *PRNP* Codon 129. *Lancet* 2009;374:2128.



**Table 2a Cases of definite and probable variant CJD shown by residence at onset (n=143) and residence at death (n=144†) in England (region & local authority)**

ENGLAND	No. resident at onset	No. resident at death	Mortality rate*	ENGLAND	No. resident at onset	No. resident at death	Mortality rate*
<b>North East</b>	<b>11</b>	<b>11</b>	<b>0.17</b>	<b>East</b>	<b>13</b>	<b>13</b>	<b>0.09</b>
Darlington UA	0	0		Luton UA	0	0	
Hartlepool UA	0	0		Peterborough UA	0	0	
Middlesbrough UA	1	1		Southend-on-Sea UA	1	1	
Redcar & Cleveland UA	1	1		Thurrock UA	0	0	
Stockton-on-Tees UA	1	1		Bedfordshire	0	0	
Durham	0	1		Cambridgeshire	1	1	
Northumberland	3	4		Essex	2	2	
Tyne & Wear	5	3		Hertfordshire	3	3	
				Norfolk	3	3	
<b>North West</b>	<b>27</b>	<b>27</b>	<b>0.15</b>	Suffolk	3	3	
Blackburn with Darwen UA	0	0		<b>London</b>	<b>20</b>	<b>18</b>	<b>0.09</b>
Blackpool UA	1	1		Inner London	7	7	
Halton UA	0	0		Outer London	13	11	
Warrington UA	2	2		<b>South East</b>	<b>23</b>	<b>20</b>	<b>0.09</b>
Cheshire	5	6		Bracknell Forest UA	1	1	
Cumbria	1	1		Brighton and Hove UA	0	0	
Greater Manchester	10	9		Isle of Wight UA	0	1	
Lancashire	4	4		Medway UA	0	1	
Merseyside	4	4		Milton Keynes UA	0	0	
<b>Yorkshire and the Humber</b>	<b>17</b>	<b>18</b>	<b>0.14</b>	Portsmouth UA	1	2	
East Riding of Yorkshire UA	1	1		Reading UA	0	0	
Kingston Upon Hull, UA	0	0		Slough UA	0	0	
North East Lincolnshire UA	1	1		Southampton UA	1	0	
North Lincolnshire UA	0	0		West Berkshire UA	0	0	
York UA	0	0		Windsor & Maidenhead UA	0	0	
North Yorkshire	4	4		Wokingham UA	0	0	
South Yorkshire	5	5		Buckinghamshire	0	1	
West Yorkshire	6	7		East Sussex	2	2	
<b>East Midlands</b>	<b>8</b>	<b>10</b>	<b>0.09</b>	Hampshire	5	2	
Derby UA	0	0		Kent	5	4	
Leicester UA	0	0		Oxfordshire	1	1	
Nottingham UA	0	0		Surrey	6	4	
Rutland UA	0	0		West Sussex	1	1	
Derbyshire	0	1		<b>South West</b>	<b>18</b>	<b>17</b>	<b>0.13</b>
Leicestershire	4	5		Bath & NE Somerset UA	0	0	
Lincolnshire	2	2		Bournemouth UA	1	1	
Northamptonshire	1	1		Bristol, City of UA	1	1	
Nottinghamshire	1	1		North Somerset UA	0	0	
<b>West Midlands</b>	<b>6</b>	<b>10</b>	<b>0.07</b>	Plymouth UA	0	0	
Herefordshire, County of UA	0	0		Poole UA	0	0	
Stoke-on-Trent UA	0	0		South Gloucestershire UA	1	0	
Telford and Wrekin UA	0	0		Swindon UA	0	0	
Shropshire	1	1		Torbay UA	0	1	
Staffordshire	0	0		Cornwall and Isles of Scilly	2	1	
Warwickshire	2	3		Devon	3	3	
West Midlands (Met County)	3	5		Dorset	0	0	
Worcestershire	0	1		Gloucestershire	2	2	
				Somerset	4	5	
				Wiltshire	4	3	
<b>TOTAL FOR ENGLAND</b>	<b>143</b>	<b>144</b>	<b>0.11</b>				

\* number of deaths/million/annum based on mid 2005 population estimates (source: ONS): 1 May 1995 to 31 Dec 2020. Postcode of residence obtained from AFD Postcode Plus.

† excludes 3 cases who died abroad.

**Table 2b Cases of definite and probable variant CJD shown by residence at onset (n=35) and residence at death (n=31): Wales, Scotland and NI**

<b>WALES†</b>	<b>No. resident at onset</b>	<b>No. resident at death</b>	<b>WALES†</b>	<b>No. resident at onset</b>	<b>No. resident at death</b>
Isle of Anglesey	0	0	Neath Port Talbot	0	0
Gwynedd	1	1	The Vale of Glamorgan	1	1
Conwy	0	0	Cardiff	0	0
Denbighshire	1	0	Bridgend	0	0
Flintshire	0	0	Rhondda, Cynon, Taff	0	0
Wrexham	0	0	Merthyr Tydfil	0	0
Powys	1	1	Caerphilly	0	0
Ceredigion	0	0	Blaenau Gwent	0	0
Pembrokeshire	2	2	Torfaen	0	0
Cardiganshire	1	1	Monmouthshire	0	0
Swansea	1	0	Newport	0	0
<b>TOTAL (MORTALITY RATE*)</b>	<b>8</b>	<b>6 (0.08)</b>	†unitary authorities		


<b>SCOTLAND†</b>	<b>No. resident at onset</b>	<b>No. resident at death</b>	<b>SCOTLAND†</b>	<b>No. resident at onset</b>	<b>No. resident at death</b>
Aberdeen City	1	1	Highland	3	2
Aberdeenshire	0	0	Inverclyde	0	0
Angus	0	0	Midlothian	0	0
Argyll & Bute	0	0	Moray	0	0
Clackmannanshire	0	0	North Ayrshire	0	0
Dumfries & Galloway	0	0	North Lanarkshire	3	3
Dundee City	0	0	Orkney Islands	1	0
East Ayrshire	1	1	Perth & Kinross	0	0
East Dunbartonshire	1	1	Renfrewshire	1	1
East Lothian	0	0	Scottish Borders	0	0
East Renfrewshire	1	1	Shetland Islands	0	0
Edinburgh, City of	2	2	South Ayrshire	1	1
Eilean Siar	0	0	South Lanarkshire	1	1
Falkirk	1	1	Stirling	0	0
Fife	2	2	West Dunbartonshire	0	0
Glasgow, City of	3	3	West Lothian	2	2
<b>TOTAL (MORTALITY RATE*)</b>	<b>24</b>	<b>22 (0.17)</b>	†council areas		

<b>N IRELAND†</b>	<b>No. resident at onset</b>	<b>No. resident at death</b>	<b>N IRELAND†</b>	<b>No. resident at onset</b>	<b>No. resident at death</b>
Antrim & Newtonabbey	1	1	Fermanagh and Omagh	0	0
Ards and North Down	0	0	Lisburn and Castlereagh	0	0
Armagh, Banbridge & Craigavon	0	0	Mid and East Antrim	0	0
Belfast	2	2	Mid Ulster	0	0
Causeway Coast and Glens	0	0	Newry, Mourne and Down	0	0
Derry City and Strabane	0	0			
<b>TOTAL (MORTALITY RATE*)</b>	<b>3</b>	<b>3 (0.07)</b>	†local government districts		

\* number of deaths/million/annum based on mid-2005 population estimates (source: ONS): 1 May 1995-31 Dec 2020.  
Postcode of residence obtained from AFD Postcode Plus.

Cases have been widely spread throughout the UK. Age- and sex- standardised incidence ratios (SIRs) based on cases' place of residence in 1991 (shortly after the time when exposure to the BSE agent is assumed to have peaked) are shown in Figure 7. There remains a relatively high incidence amongst those who lived in the north (Scotland, North East, North West, Yorkshire & Humberside; 16.9 million people, 74 vCJD cases) compared to the south (Wales, East Midlands, West Midlands, South West, South East, London, East of England; 31.7 million people, 100 vCJD cases) of Great Britain in 1991.<sup>2</sup> The rate ratio controlling for age and sex is 1.39 (95% CI 1.03-1.88), ie individuals living in the “North” in 1991 are about one and a half times more likely to have developed vCJD than individuals who were living in the “South” in 1991.

**Figure 7** Standardised variant CJD incidence ratios (SIRs) up to 31<sup>st</sup> December 2020, by region of residence on 1st January 1991




<sup>2</sup> Cousens S, Smith PG, Ward H, Everington D, Knight RSG, Zeidler M, Stewart G, Smith-Bathgate EAB, Macleod MA, Mackenzie J, Will RG. Geographical distribution of variant Creutzfeldt-Jakob disease in Great Britain, 1994-2000. *Lancet* 2001; 357: 1002-1007.

## 2.4 Iatrogenic Creutzfeldt-Jakob disease

Since 1970, up to 31st December 2020, 89 cases of CJD attributable to iatrogenic exposure have been identified, 8 in individuals receiving dura mater implants, 80 in individuals who had received human-derived growth hormone (hGH) and one in a recipient of human gonadotrophin (hGN) who was treated in Australia. All of these individuals have died (Figure 8). The median age at death of the hGH/hGN group was 35 years (with a range of 20-51 years) and for the dura mater cases 46 years (range 27-78 years).

**Figure 8** Deaths from iatrogenic CJD, 1979-2020



The first identified iatrogenic case was a dura mater recipient who died in 1979. The first hGH-related death occurred in 1985. Since 1985 in the UK, human pituitary-derived hormones have been replaced by synthetic preparations. Details of the UK human pituitary-derived hormone cases, with a discussion of the incubation periods, were published in 2003.<sup>3</sup> A study of the accumulated UK experience with dura mater-related CJD, including incubation periods, was undertaken and the results published in 2006.<sup>4</sup>

Iatrogenic transmission of CJD/vCJD is also studied by the Unit through the identification and investigation of surgical or other links between cases. The Unit continues to collect risk factor information for all suspect cases of human prion diseases referred to the Unit as part of its core work.

## 2.5 Transfusion Medicine Epidemiology Review

The Transfusion Medicine Epidemiology Review (TMER) is a collaborative project between the UK NCJDRSU and UK Blood Services (UKBS). The main purpose is to investigate whether there is any evidence that CJD or vCJD may have been transmitted via the blood supply. Cases (definite and probable) are notified to the UKBS by NCJDRSU; a search establishes whether any have acted as donors or received blood transfusions. Donation/transfusion records are checked and all

<sup>3</sup> Swerdlow AJ, Higgins CD, Adlard P, Jones ME, Preece MA. Creutzfeldt-Jakob disease in United Kingdom patients treated with human pituitary growth hormone. *Neurology* 2003; 61: 783-91.

<sup>4</sup> Heath CA, Barker RA, Esmonde TFG, Harvey P, Trend P, Head MW, Smith C, Bell JE, Ironside JW, Will RG, Knight RSG. Dura mater-associated Creutzfeldt-Jakob disease: experience from surveillance in the UK. *JNNP* 2006; 77: 880-2.

components traced through hospital records. Details of all identified recipients/donors are forwarded to NCJDRSU for subsequent checking to ensure none appear on the NCJDRSU database as a case of CJD.

Results from the vCJD arm of the project identified four instances of probable transfusion transmitted infection in 3 cases of vCJD and pre-clinical infection in a recipient with post-mortem confirmation of abnormal prion protein deposition in the spleen (all previously reported<sup>5678</sup>). There have been no new cases of transfusion-associated vCJD since 2007.

Results from all other types of CJD included in the project have not so far shown any evidence of transfusion transmission<sup>9</sup>. This includes, to date, 391 blood component recipients identified from 55 sporadic CJD (sCJD) cases who were donors. None have been identified as CJD cases on the NCJDRSU database and death certificates from the 196 recipients who have subsequently died did not reveal CJD as a cause of death. Twenty-nine sCJD cases with a history of blood component transfusion were traced by the blood services from which 292 donors were identified. None of these donors have been identified as CJD cases on the NCJDRSU database and death certificates from the 11 donors who have subsequently died did not reveal CJD as a cause of death.

(External collaborators on this project: Dr H Harvala Simmonds, Ms C Reynolds, Ms T Yawitch).

## **2.6 Study of Progressive Intellectual & Neurological Deterioration (PIND)**

The aim of this project is to use the mechanism of the British Paediatric Surveillance Unit to identify all cases of progressive intellectual and neurological deterioration in children in the UK, particularly those with features suggestive of vCJD. All cases are discussed and allocated to a diagnostic category by an Expert Neurological Advisory Group made up of consultants who have specialised knowledge of paediatric neurology, neurogenetics and metabolic disease, together with representation from the National CJD Research & Surveillance Unit.<sup>101112</sup>

As of 31<sup>st</sup> December 2020, after nearly 24 years of surveillance, 4774 patients with suspected PIND had been reported. There have been six cases of vCJD in children aged 15 years or less: four definite and two probable. Three were reported in 1999, one in 2000 and two in mid-2001. The youngest UK case of vCJD identified to date was aged 12 at onset. A total of 2076 cases had a confirmed underlying cause other than vCJD, being categorised into over 220 known neurodegenerative diseases.

(External collaborators on this project: Dr C Verity, Mrs E Baker, Ms AM Winstone, Ms P Maunder)

<sup>5</sup> Llewelyn CA, Hewitt PA, Knight RSG, Amar K, Cousens S, Mackenzie J, Will RG. Possible transmission of variant Creutzfeldt-Jakob disease by blood transfusion. *Lancet* 2004; 363: 417-421.

<sup>6</sup> Peden AH, Head MW, Ritchie DL, Bell JE, Ironside JW. Preclinical vCJD after blood transfusion in a PRNP Codon 129 heterozygous patient. *Lancet* 2004 364: 527-529.

<sup>7</sup> Wroe SJ, Pal S, Siddique D, Hyare H, Macfarlane R, Joiner S, Linehan JM, Brandner S, Wadsworth JD, Hewitt P, Collinge J. Clinical presentation and pre-mortem diagnosis of variant Creutzfeldt-Jakob disease associated with blood transfusion: a case report. *Lancet* 2006; 368: 2061-2067.

<sup>8</sup> Health Protection Agency. Fourth case of transfusion-associated variant-CJD. *Health Protection Report* 2007;1(3):

<sup>9</sup> Urwin PJ, Mackenzie JM, Llewelyn CA, Will RG, Hewitt PE. Creutzfeldt-Jakob disease and blood transfusion: updated results of the UK Transfusion Medicine Epidemiology Review Study. *Vox Sang* 2016; 110: 310-316.

<sup>10</sup> Verity CM, Nicoll A, Will RG, Devereux G, Stellitano L. Variant Creutzfeldt-Jakob disease in UK children: a national surveillance study. *Lancet* 2000; 356: 1224-1227.

<sup>11</sup> Devereux G, Stellitano L, Verity CM, Nicoll A, Will RG, Rogers P. Variations in neurodegenerative disease across the UK: findings from the national study of Progressive Intellectual and Neurological Deterioration (PIND). *Arch Dis Child* 2004; 89: 8-12.

<sup>12</sup> Verity C, Winstone AM, Stellitano L, Will R, Nicoll A. The epidemiology of progressive intellectual and neurological deterioration in childhood. *Arch Dis Child* 2010; 95:361-364 deterioration in childhood. *Arch Dis Child* 2010; 95:361-364.

## **2.7 Surveillance of potential occupational exposure to CJD**

Public Health England in collaboration with NCJDRSU have set up an occupational surveillance study with two parts: 1) a registry for the prospective long term monitoring of healthcare and laboratory workers who have incurred occupational exposures to prion diseases and 2) the retrospective review of possible occupational exposures of CJD cases who have been healthcare or laboratory workers. By the end of 2020, 2 healthcare workers and one laboratory worker had reported prion-disease exposures as a result of needle stick/sharps injuries. None have subsequently developed prion disease. Retrospective investigations of possible occupational exposures of CJD cases in the UK continues to be undertaken to determine if any exposure to prion disease occurred – there is no evidence to indicate the occurrence of occupational exposure to the prion agent.<sup>13 14</sup>

(External collaborators on this project: K Sinka)

## **2.8 Prion surveillance in primary immunodeficiency patients**

The study began in 2006 and aims to identify whether there is evidence of abnormal prion protein/vCJD in the blood and/or body tissues of primary immunodeficiency patients exposed to UK sourced immunoglobulin between 1996 and 2000.

By 31<sup>st</sup> December 2020, a total of 80 patients registered in 17 immunology centres across Great Britain had participated in the study. Of these, 18 had died with a further 9 lost to follow up, leaving 53 participants registered over 12 sites. Participants have been followed up for approximately 1566 person-years following first exposure to UK-sourced immunoglobulin. In this time no participants have shown any clinical or pathological features suggestive of vCJD or evidence of abnormal prion protein in tissues tested.

(External collaborators on this project: M Turner, R McNairney, M Helbert, M Buckland, J Cooper, R Herriott, A Huissoon, M Gompels, S Jolles, C Chopra, G Hayman, P Wood, M Browning, T Garcez, A Herwadkar, D Lowe, M Thomas, C Bethune, S Goddard)

## **2.9 Enhanced surveillance of individuals identified as at increased risk of CJD**

The potential for secondary transmission of CJD has led to collaborative studies undertaken between the UK Haemophilia Centre Doctors Organisation, Institute of Child Health (London), NHS Blood and Transplant, National Prion Clinic, Public Health England and Health Protection Scotland aimed at identifying whether there is evidence of clinical or sub-clinical infection in those judged to be at increased risk of CJD, such evidence is investigated through review of clinical records and medical histories, and through post-mortem investigations<sup>15</sup>.

As at 31<sup>st</sup> December 2020, three cases of vCJD and one asymptomatic infection had been identified in recipients of blood from donors who later developed vCJD (see section 2.5 TMER) and one asymptomatic infection in a bleeding-disorder patient who received UK sourced plasma products.

<sup>13</sup> Thorpe J, Mackenzie J, Molesworth A, Sinka K, Will R. Occupational exposures to prion diseases in healthcare and laboratory workers. Poster presentation at Prion 2012, 9-12 May, Amsterdam.

<sup>14</sup> Mackenzie JM, Urwin P, Mackenzie G, Knight RSG, Will RG, Molesworth AM. Occupations of cases of vCJD in the UK. Poster presentation at Prion 2017, 23-26 May, Edinburgh.

<sup>15</sup> Public Health England. Creutzfeldt-Jakob Disease (CJD) Biannual Update (February 2018). Health Protection Report, Vol 12, Number 5, 9 February 2018.

There have been no occurrences/diagnoses of CJD in individuals at risk through surgical exposures. Please see section 2.4 for figures relating to those at risk following treatment with pituitary derived hormones.

(External collaborators on this project: (H Ward, K Sinka, S Mead)

## LABORATORY ACTIVITIES

Laboratory investigations are part of the internationally-agreed diagnostic criteria for CJD, both during life (CSF protein analysis, PrP genetic studies, brain biopsy neuropathology and prion protein studies) and post-mortem (autopsy neuropathology and prion protein studies). The NCJDRSU has facilities to perform all of these investigations, which aid the timely and accurate diagnosis of all forms of CJD and are essential for surveillance purposes.

### ***3.1 Neuropathology – Statement of Progress and Surveillance Activities***

The neuropathology laboratory in the NCJDRSU continues to maintain its diagnostic and research activities, with most of the cases investigated referred from other centres across the UK (see Table 3). The laboratory maintains close links with other neuropathology centres across the UK and overseas with scientific, medical, technical advice over the past year. The laboratory has continued to maintain an active research programme both in-house and by collaboration with other research centres in UK, Europe and across the world and provides tissues to researchers through the CJD Brain and Tissue Bank, which is part of the MRC-funded Edinburgh Brain Bank.

In the contemporary referrals for 2020, the numbers of cases diagnosed as prion diseases was lower than those in the previous year; the numbers of cases in which there was no evidence of CJD or an alternative diagnosis was also lower than the previous year. No cases of vCJD were identified in the UK and none were referred from outside the UK.

In addition to the UK CJD surveillance work, the neuropathology laboratory is involved in a number of collaborative research and surveillance studies in relation to neuropathological diagnosis of CJD and other human prion diseases.

The laboratory and its staff continue to participate in a range of EQA activities related to both technical and diagnostic neuropathology. As before, the laboratory continues to act as a source of information to a wide range of professionals involved in health and safety issues relating to CJD. We are most grateful to all neuropathologists, general pathologists and their technical, secretarial and autopsy room staff for their continuing support of the NCJDRSU. We are also grateful to the relatives of patients with CJD for allowing us to study this group of devastating disorders.



**Table 3 Breakdown of Laboratory Activities:  
Period 1<sup>st</sup> January 2020– 31<sup>st</sup> December 2020**

	2020	2019
<b>REFERRED CASES (UK)</b>		
Sporadic CJD	12	15 <sup>1</sup>
Genetic CJD	0	1
Variant CJD	0	
Iatrogenic CJD (GHT)	0	0
Iatrogenic CJD (Lyodura)	0	0
Gerstmann-Straussler-Scheinker Syndrome	0	0
Fatal Familial Insomnia	0	
Variably protease sensitive prionopathy	2	1
No evidence of CJD	2	1
Alzheimer's disease		0
Lewy Body disease	0	0
Other forms of brain disease <sup>2</sup>	2	0
<b>REFERRED CASES (EU)</b>		0
Sporadic CJD	0	0
Familial CJD	0	1
GSS	1	0
Fatal Familial Insomnia		
No evidence of CJD	0	0
<b>REFERRED CASES (ROW)</b>		
CJD, presumed sporadic <sup>1</sup>	0	0
Other forms of brain disease <sup>2</sup>	0	0
		1
<b>TOTAL NUMBER OF CASES</b>	<b>35</b>	<b>48</b>

## NOTES

- 1 Confirmed CJD that, in the absence of genetic testing for an inherited mutation, is presumed sporadic in origin.
- 2 Other (2020) Meningioma n=1; B-cell lymphoma n=1; Hippocampal Sclerosis n=1; Small vessel disease n=2; Global Ischaemic injury n=1; FTLD n=1; Familial amyloid angiopathy n=1

## Abbreviations:

GHT Growth Hormone Therapy  
ROW Rest of World  
EU European Union

## 3.2 Protein Biochemistry Laboratory

### Prion protein detection and typing

Prion protein typing is carried out as a routine diagnostic test on all suspected cases of CJD from which frozen brain tissue is received by the NCJDRSU. Small quantities of cerebral cortex or cerebellum are homogenised, treated with protease and the size and relative abundance of the protease resistant prion protein (PrP<sup>res</sup>) fragments determined by Western blot analysis. The recognised PrP<sup>res</sup> types, their nomenclature and their association with different human prion diseases are shown in Figure 9 and described in the accompanying legend. In cases from which only peripheral tissues are available (such as those in which diagnostic tonsil biopsy is performed), or in cases in which the patient is thought to have been at risk of developing CJD due to potential iatrogenic exposure and is enrolled in a UK prion screening study, a modified Western blot procedure is used which employs centrifugal concentration or sodium phosphotungstic acid precipitation to enrich for PrP<sup>res</sup> and increase the sensitivity of the test.

Figure 9




Figure 9 shows the diagrammatic representation of the main protease resistant prion protein (PrP<sup>res</sup>) types found in the human prion disease brain as determined by proteinase K digestion and Western blot analysis. The classification of the banding pattern has two components, one numerical depending on the migration of the bands and the other alphabetical depending on their relative abundance. The pattern is termed type 1 if the non-glycosylated (bottom) band is ~21kDa, type 2 if the non-glycosylated band is ~19kDa or type 1+2 if both bands are found. In cases and samples in which both types are present but one type predominates the less abundant type is placed in parentheses [ie type 1(+2) or type 2(+1)]. The pattern is given the suffix A if the middle or bottom (mono-, or non-glycosylated) bands predominate, B if the top (di-glycosylated) band predominates or A/B if the glycosylated bands (middle and top) predominate at the expense of the non-glycosylated (bottom) band. A pattern dominated by a low molecular mass unglycosylated band is here termed 8kDa. The faint ladder of bands that sometimes accompanies the 8kDa band is shown in grey. Types 1A, 2A, 1+2(A) are characteristic of sporadic and iatrogenic CJD. Type 2B is associated with variant CJD and is a consistent feature present in all cases so far examined. However, a protein isotype resembling type 2B can also be found in cases of FFI and fCJDE200K. Types 1B, 1A/B and 2A/B are often found in genetic CJD, GSS and FFI. The 8kDa pattern is characteristic of some cases of GSS and of VPSPr. VPSPr can present with multiple isotypes, one of which includes an intermediate band also associated with iatrogenic CJD.

### UK Referrals

A total of 16 UK cases with frozen tissue were received and analysed in 2020. The results of the analysis were as follows:

**Table 4 Breakdown of cases analysed in 2020**

Diagnosis	Type	PrP <sup>res</sup> +ve CNS
CJD	Sporadic	10
VPSPr		1
Alternative final diagnosis or not determined		5

Further sub-classification by PrP<sup>res</sup> type and PRNP genotype yields the following results:

**Table 5 PrP<sup>res</sup> type / PRNP genotype breakdown of CJD cases analysed in 2020**

Diagnosis	PRNP genotype	Type 1A	Type 2A	Type LMW <sub>t</sub> (8Kda)
Sporadic CJD	MM	5	1	
	MV		2	
	VV		2	
VPSPr	VV			1 <sup>1</sup>

<sup>1</sup> Includes one case with type 8kDa plus a minority type 2A

### ***Non-UK referrals***

A total of 7 non-UK cases with frozen tissue were received and analysed in 2020. The results of the analysis were as follows:

**Table 6 Breakdown of cases analysed in 2020**

Diagnosis	Type	PrP <sup>res</sup> +ve CNS
CJD	Sporadic	6
GSS		1

There were 7 non-UK referrals for Western blot analysis on frozen tissue, all were sCJD or GSS. Four cases came from Sweden, one from Ireland and two from New Zealand.

**Table 7 PrP<sup>res</sup> type / PRNP genotype breakdown of cases analysed in 2020**

Diagnosis	PRNP genotype	Type 1A	Type 2A	Type LMW <sub>t</sub> (8Kda)
Sporadic CJD	MM	2	2	
	MV		1	
	VV		1	
GSS (P102L)	MM			1

### **3.3 Brain banking activities**

The neuropathology laboratory houses the CJD Brain and Tissue Bank, which is part of the Edinburgh Brain and Tissue Bank, directed by Professor Colin Smith. The CJD Brain and Tissue Bank was used extensively in 2020 for diagnostic and collaborative research purposes with colleagues in the UK and overseas. The Edinburgh Brain and Tissue Banks are part of the MRC Network of UK Brain Banks, which works to strengthen banking activities and ensure uniform high standards of operation. The Bank has a website, on which further details are available including instructions on how to request tissue samples for research (<http://www.ed.ac.uk/clinical-brain-sciences/research/edinburgh-brain-and-tissue-bank>). The activities of the Bank comply with current

guidelines from the Royal College of Pathologists, the Human Tissue (Scotland) Act 2006 and the Human Tissue Act 2004.

### 3.4 Molecular Genetics

#### Genetic CJD

One hundred and eighty-two cases of genetic CJD (excluding cases of GSS) have been identified since 1970 by the NCJDRSU (these data are incomplete as formal investigation of genetic CJD in the UK is undertaken by the National Prion Clinic in London). Of the 182 cases, 161 were resident in England, 11 were resident in Wales, 3 were resident in Northern Ireland and 7 were resident in Scotland. Twelve cases were still alive as at 31<sup>st</sup> December 2020 and one has been lost of follow-up after moving out of the UK after diagnosis. Seventy-four of the cases had insertions in the coding region of the PrP gene, 60 carried the mutation at codon E200K, 19 at codon D178N, 4 at codon V210I, one at codon D167G, 2 at codon V163STOP, one at codon G54S, one at codon E211Q, 3 at codon E196K and one at codon P105S. The remaining 16 were identified as genetic on the basis of relatives known to have had CJD. The mean age at death was 56½ and median was 56 years (range 29-95 years).

#### PRNP Codon 129 distribution in sporadic CJD

The distribution of PRNP Codon 129 genotypes in sCJD has been analysed since the inception of the Unit in 1990. The overall distribution of PRNP Codon 129 genotypes in sCJD is 61% MM, 20% MV, 19% VV (see Table 8). There appears to be evidence ( $p<0.05$ ) of a change in the PRNP Codon 129 distribution in sCJD between the first and subsequent periods. The explanation for this remains unclear. It should be noted that not all cases are genotyped (data available on 62%) and, therefore, changes in PRNP Codon 129 distribution may reflect changes in the way in which cases are selected for analysis.

**Table 8 PRNP Codon 129 genotypes of cases of sporadic CJD in the UK, 1990-2020**

Deaths from sCJD	Percentage tested	MM (%)	MV (%)	VV (%)
Deaths from 1 January 1990 – 31 December 1999	63%	200 (70)	43 (15)	43 (15)
Deaths from 1 January 2000 – 31 December 2009	63%	255 (59)	90 (21)	87 (20)
Deaths from 1 January 2000 – 31 December 2009	63%	436 (58)	159 (21)	157 (21)
Total	63%	891 (61)	292(20)	287 (19)
<b>Genotype distribution for the normal population<sup>16</sup></b>		(44)	(45)	(11)

#### PRNP Codon 129 distribution in variant CJD

In clinical cases for whom genetic data are available ( $n=161$ , 90%), 160 were methionine homozygotes at PRNP Codon 129 of the PrP gene and one case was heterozygous at PRNP Codon 129 of the PrP gene.

The genetic laboratory undertakes genetic analysis on a national and international basis.

<sup>16</sup> Bishop et al. PRNP variation in UK sporadic and variant Creutzfeldt-Jakob disease highlights genetic risk factors and a novel non-synonymous polymorphism. BMC Medical Genetics 2009;10:146-155.

### 3.5 CSF RT-QuIC, 14-3-3 and other brain specific proteins

#### Introduction

During the period January-December 2020, the laboratory received 256 cerebrospinal fluid (CSF) from suspected CJD patients residing in the UK, 76 samples from suspected CJD patients residing outwith the UK and 22 from young onset dementia patients (Table 9).

**Table 9 Origin of CSF samples sent to the NCJDRSU for CSF 14-3-3 analysis from January 2020 – December 2020**

Patient Group	Number CSF samples received
Young Onset Dementia	22
Suspected CJD (non-UK)	76
Suspected CJD (UK)	256
Total number	354

Results of 14-3-3 and RT-QuIC analysis on the 256 cases of suspected CJD in the UK are shown in Table 10.

**Table 10 CSF 14-3-3 and RT-QuIC results in 256 CSF samples from suspected CJD cases in the UK**

Patient Group (n)	14-3-3 number positive/total number analysed (% positive)	RT-QuIC number positive/total number analysed (% positive)
Neuropathologically confirmed sporadic CJD (3)	1/3 (33%)	2/3 (67%)
Probable sporadic CJD (103*)	50/95* (53%)	98/103 (95%)
Possible sporadic CJD (1)	0/1	0/1
Neuropathologically confirmed Variably Protease Sensitive Prionopathy (1)	0/1	0/1
Probable genetic D178N 129-M CJD (1)	0/1	0/1
Not CJD (147)	6/142** (4%)	0/147 (0%)

\*Eight CSF samples were insufficient for both CSF 14-3-3 and RT-QuIC analysis, so CSF RT-QuIC was prioritized.

\*\* Five CSF samples were insufficient for both CSF 14-3-3 and RT-QuIC analysis, so CSF RT-QuIC was prioritized.

We received CSF samples from 106 of the 130 (82%) neuropathologically confirmed or probable sporadic CJD cases referred to the NCJDRSU during January – December 2020. Despite the two national lockdowns introduced by the national government in 2020, there was no significant reduction in the numbers of CSF samples received or analysed by the NCJDRSU.

Details of the investigations in the 6 sporadic CJD patients who had a negative RT-QuIC result are given in Table 11.

**Table 11** Cases of sporadic CJD with negative RT-QuIC result

<b>Id</b>	<b>CSF 14-3-3</b>	<b>CSF S-100b (ng/ml)*</b>	<b><i>PRNP</i> Codon 129 genotype</b>	<b>PrP<sup>Sc</sup> isotype</b>	<b>Disease Duration (months)</b>	<b>Cortical ribboning Y/N</b>	<b>Basal Ganglia Changes Y/N</b>	<b>Comments (MRI)</b>
1	Negative	1.10	MM	2A	13	Y	Y	Positive scan
2	Weak Positive	0.29				Y	Y	Positive scan
3	Positive	0.53	VV	nd	3	N	Y	Atypical caudate change
4	Weak Positive	0.52	MM	nd	nd	Y	Y	Positive scan
5	Negative	0.41		nd		Y	Y	Positive scan
6	Positive	0.53	MV	nd	Still alive	N	N	Positive scan

\*Reference range <0.41 ng/ml; nd: not done

## **NATIONAL CJD CARE TEAM**

**E**stablished by the Department of Health and Social Care, the National CJD Care Team is based within the National CJD Research & Surveillance Unit and was formed in order to optimise the care of patients suffering from all forms of CJD. The national care co-ordinator post was established in February 2000, and the National CJD Care Team formed in September 2001. They provide an NHS service to England, Wales, Scotland and Northern Ireland. The present team consists of two care co-ordinators who are senior nurses with the support of a consultant neurologist, research registrars and a secretary.

During the period 1<sup>st</sup> January to 31<sup>st</sup> December 2020, 137 new referrals were made to the Care Team. All new referrals with suspected CJD are assessed by a care co-ordinator in person whenever possible. During the pandemic this was limited to telehealth and telephone assessments, meetings and consultations. This was so successful that going forward the service will be a combination of both systems.

The co-ordinators work closely with family and local healthcare professionals in assisting with discharge and care planning. Care co-ordinators are able to meet with patients, families and local professionals depending on individual need. They provide valuable expertise in nursing patients with CJD and in anticipation and prevention of problems that may arise. Care co-ordinators are available to provide advice and education on diagnosis, prognosis, discharge planning, symptom management, infection control and any other questions in relation to the care of patients with CJD. Contact is maintained in person, by telephone, email or telehealth.

The care co-ordinators play a pivotal role in shaping the patient and family experience of CJD. They work with that family to identify preferences for information and how they are going to get that information. The care co-ordinators often have to clarify misunderstandings and explain complex terminology. They provide information about the next steps in the care pathway which will include the expected course of the disease and its symptomatic treatment with all its potential consequences for the patient and family.

The co-ordinators provide information and opportunities for discussion as families and sometimes patients address complex issues like DNACPR, informed consent, advanced directives, feeding concerns, end of life decisions and the health care team. They act as an advocate for the patient and family and often as an intermediary between the family and the health care team. All this whilst supporting the family emotionally through a devastating diagnosis that has no answers as to why them and has often began very quickly and progressed rapidly.

The National CJD Care Team works in close liaison with NHS England and provides access to the CJD Care Package. This provides additional funding to supplement local authorities with the care of patients suffering from all forms of CJD. Care packages are flexible and can change quickly to meet the rapidly changing needs of patients. The aim is to provide a package of care that will meet the

needs both for the patient and their family in a timely manner. In addition to collaborations with national organisations in the United Kingdom, the Care Team liaises closely with international organisations, including the Australian and American CJD Support Groups and is an Official Friend of the CJD International Support Alliance.



## PUBLICATIONS IN 2020

1. Cassard H, Huor A, Espinosa JC, Douet JY, Lugan S, Aron N, Vilette D, Delisle MB, Marín-Moreno A, Peran P, Beringue V, Torres JM, Ironside JW, Andreoletti O. Prions from sporadic Creutzfeldt-Jakob Disease patients propagate as strain mixtures. *mBio*. 2020; 11(3): e00393-20.
2. Gill ON, Spencer Y, Richard-Loendt A, Kelly C, Brown D, Sinka K, Andrews N, Dabaghian R, Simmons M, Edwards P, Bellerby P, Everest DJ, McCall M, McCardle LM, Linehan J, Mead S, Hilton DA, Ironside JW, Brandner S. Prevalence in Britain of abnormal prion protein in human appendices before and after exposure to the cattle BSE epizootic. *Acta Neuropathol*. 2020; 139(6): 965-976.
3. Han JY, Jang HS, Green AJE, Choi YP. RT-QuIC-based detection of alpha-synuclein seeding activity in brains of dementia with Lewy Body patients and of a transgenic mouse model of synucleinopathy. *Prion*. 2020; 14(1): 88-94.
4. Jones E, Hummerich H, Viré E, Uphill J, Dimitriadis A, Speedy H, Campbell T, Norsworthy P, Quinn L, Whitfield J, Linehan J, Jaunmuktane Z, Brandner S, Jat P, Nihat A, How Mok T, Ahmed P, Collins S, Stehmann C, Sarros S, Kovacs GG, Geschwind MD, Golubjatnikov A, Frontzek K, Budka H, Aguzzi A, Karamujić-Čomić H, van der Lee SJ, Ibrahim-Verbaas CA, van Duijn CM, Sikorska B, Golanska E, Liberski PP, Calero M, Calero O, Sanchez-Juan P, Salas A, Martín-Torres F, Bouaziz-Amar E, Haik S, Laplanche JL, Brandel JP, Amouyel P, Lambert JC, Parchi P, Bartoletti-Stella A, Capellari S, Pileggi A, Ladogana A, Pocchiari M, Aneli S, Matullo G, Knight R, Zafar S, Zerr I, Booth S, Coulthart MB, Jansen GH, Glisic K, Blevins J, Gambetti P, Safar J, Appleby B, Collinge J, Mead S. Identification of novel risk loci and causal insights for sporadic Creutzfeldt-Jakob disease: a genome-wide association study. *Lancet Neurol*. 2020; 19(10): 840-848.
5. Knight R. Clinical diagnosis of human prion disease. *Prog Mol Biol Transl Sci*. 2020; 175: 1-18.
6. Lauwers E, Lalli G, Brandner S, Collinge J, Compennolle V, Duyckaerts C, Edgren G, Haik S, Hardy J, Helmy A, Iverson AJ, Jaunmuktane Z, Jucker M, Knight R, Lemmens R, Lin IC, Love S, Mead S, Perry VH, Pickett J, Poppy G, Radford SE, Rousseau F, Routledge C, Schiavo G, Schymkowitz J, Selkoe DJ, Smith C, Thal DR, Theys T, Tiberghien P, van den Burg P, Vandekerckhove P, Walton C, Zaaier HL, Zetterberg H, De Strooper B. Potential human transmission of amyloid  $\beta$  pathology: surveillance and risks. *Lancet Neurol*. 2020; 19(10): 872-878. Epub 2020 Sep 16. Erratum in: *Lancet Neurol*. 2020;19(11): e10.

7. Orrú CD, Groveman BR, Foutz A, Bongianni M, Cardone F, McKenzie N, Culeux A, Poleggi A, Grznarova K, Perra D, Fiorini M, Liu X, Ladogana A, Sbriccoli M, Hughson AG, Haik S, Green AJ, Geschwind MD, Pocchiari M, Safar JG, Zanusso G, Caughey B. Ring trial of 2nd generation RT-QuIC diagnostic tests for sporadic CJD. *Ann Clin Transl Neurol.* 2020; 7(11): 2262-2271.
8. Verity C, Baker E, Maunder P, Pal S, Winstone AM. Differential diagnosis of progressive intellectual and neurological deterioration in children. *Dev Med Child Neurol.* 2021; 63(3): 287-294. [Epub 2020 Sep 24.](#)
9. Ward A, Hollister JR, McNally K, Ritchie DL, Zanusso G, Priola SA. Transmission characteristics of heterozygous cases of Creutzfeldt-Jakob disease with variable abnormal prion protein allotypes. *Acta Neuropathol Commun.* 2020; 8(1): 83.
10. Watson N, Kurudzhu H, Green A, Summers D, Smith C, Pal S. Application of telehealth for comprehensive Creutzfeldt-Jakob disease surveillance in the United Kingdom. *J Neurol Sci.* 2021; 420: 117221. [Epub 2020 Nov 7.](#)

***Staff based at the National CJD Research & Surveillance Unit, Edinburgh in 2020***

Dr C Smith	Director, NCJDRSU, Professor of Clinical Neuropathology
Dr AM Molesworth	Deputy Director, NCJDRSU, Senior Epidemiologist
Dr A Green	Reader (CSF analysis)
Dr S Pal	Senior Clinical Lecturer in Neurology and Honorary Consultant Neurologist
Dr M Barria	Group Leader/Research Fellow
Professor RSG Knight	Consultant Neurologist, Professor of Clinical Neurology
Dr D Summers	Consultant Neuroradiologist
Ms J Ewins	Unit Manager
Dr L Kanguru	Senior Epidemiologist
Dr N Watson	Clinical Research Fellow
Dr H Kurudzhu	Clinical Research Fellow
Dr J Kirby	Clinical Research Fellow
Dr A Peden	Postdoctoral Research Fellow
Ms J Mackenzie	Surveillance Co-ordinator
Ms T Lindsay	European Study Co-ordinator
Mrs B Smith-Bathgate	National Care Co-ordinator and Senior Nurse
Ms M Leitch	National Care Co-ordinator and Senior Nurse
Mr N Attwood	Database Manager
Dr D Ritchie	Postdoctoral Research Fellow
Dr N McKenzie	Postdoctoral Research Fellow
Mr J Alibhai	Postdoctoral Research Fellow
Dr S Suleiman	Postdoctoral Research Scientist
Mr G Piconi	Postdoctoral Research Scientist
Ms S Lowrie	Senior Biomedical Scientist
Mrs M Le Grice	Senior Biomedical Scientist
Mrs M Andrews	Senior Biomedical Scientist
Ms H Yull	Senior Research Technician
Mr G Fairfoul	Research Technician
Ms K Burns	CSF Laboratory Technical Officer
Ms L Banks	Research Technician
Ms A Chong	Research Technical Officer
Mrs Elaine Lord	Senior Administrative Manager
Mrs F Frame	Secretariat
Mrs C Donaldson	Secretariat/Data Handler
Ms L Bond	Secretariat
Mrs K Karekwaivanane	Research Nurse
Dr S Cudmore	Data Analyst
Ms C Wardhaugh	Research Assistant
Mr F Brydon	Research Technical Officer